



Technical Data

HiEncap™ Brain Heart Infusion Broth

EC210CCL

Brain Heart Infusion Broth is employed for the propagation of fastidious pathogenic cocci and other organisms associated with blood culture work and allied pathological investigations.

Composition**

Ingredients	Gms / Litre
Calf brain, infusion from	200.000
Beef heart, infusion from	250.000
Proteose peptone	10.000
Dextrose	2.000
Sodium chloride	5.000
Disodium phosphate	2.500
Final pH (at 25°C)	7.4±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Each capsule contains 9.25 grams. Suspend 1 capsule in 250 ml (4 capsules in 1000 ml) distilled or purified water. Heat to boiling to dissolve the medium completely. Dispense into bottles or tubes and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. For best results, the medium should be used on the day it is prepared, otherwise, it should be boiled or steamed for a few minutes and then cooled before use.

Principle And Interpretation

Brain Heart Infusion Medium is useful for cultivating a wide variety of microorganisms since it is a highly nutritive medium. It is also used to prepare the inocula for antimicrobial susceptibility testing. Brain Heart Infusion Broth is a modification of the original formulation of Rosenow, where he added pieces of brain tissues to dextrose broth (1). Brain Heart Infusion Broth is also the preferred medium for anaerobic bacteria, yeasts and moulds (2-4). This medium is nutritious and well buffered to support the growth of wide variety of organisms (2, 5, 6). With the addition of 10% defibrinated sheep blood, it is useful for isolation and cultivation of *Histoplasma capsulatum* (7) and other fungi. For selective isolation of fungi, addition of gentamicin and/or chloramphenicol is recommended (8).

Proteose peptone and infusions (calf brain and beef heart) serve as sources of carbon, nitrogen, essential growth factors, amino acids and vitamins. Dextrose serves as a source of energy. Disodium phosphate helps in maintaining the buffering action of the medium whereas sodium chloride maintains the osmotic equilibrium of the medium.

Quality Control

Appearance

Gelatin capsule containing cream to yellow coloured granular media

Colour and Clarity of prepared medium

Light to medium amber coloured, clear solution without any precipitate

Quantity

Each capsule contains 9.25 gms of medium sufficient for 250 ml media.

Reaction

Reaction of 3.7% w/v aqueous solution at 25°C. pH : 7.4±0.2

pH

7.20-7.60

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 24-48 hours.

Cultural Response

Organism	Inoculum (CFU)	Growth
Cultural Response		
<i>Enterococcus faecalis</i> ATCC 50-100 29212		good-luxuriant
<i>Neisseria meningitidis</i> ATCC 50-100 13090		good-luxuriant
<i>Streptococcus pneumoniae</i> ATCC 50-100 6303		good-luxuriant
<i>Streptococcus pyogenes</i> ATCC 50-100 19615		good-luxuriant
<i>Candida albicans</i> ATCC 50-100 10231		good-luxuriant
<i>Staphylococcus aureus</i> ATCC 50-100 25923		good-luxuriant

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

Reference

- Rosenow, 1919, J. Dental Research, 1:205.
- MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.
- Atlas R. M., 1993, Handbook of Microbiological Media, 147-153, CRC Press, Boca Raton, FL.
- Downes F. P. and Ito K., (Eds.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., APHA, Washington, D.C.
- Roseburg T. et al, 1944, J. Inf. Dis., 74:131
- Conant N. F., 1950, Diagnostic Procedures and Reagents, 3rd Ed., APHA Inc., New York
- Howard B., Keiser J. F., Weissfeld A. et al, 1994, Clinical and Pathogenic Microbiology, 2nd Ed., Mosby Co.
- Murray P. R., Baron J. H., Pfaller M. A., Jorgensen J. H. and Tenover F. C., (Eds.), 2003, Manual of Clinical Microbiology, 8th Ed., American Society for Microbiology, Washington, D.C.

Revision : 00 / 2014



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.