

Oxytetra Glucose Yeast Agar Base (OGYE Agar Base), Granulated GM639I

Oxytetra Glucose Yeast Agar Base (OGYE Agar Base), granulated is recommended for isolation and enumeration of yeasts and moulds from milk and milk products. The composition and performance criteria of this medium are as per the specifications laid down in ISO 1992, ISO/DIS 6611.

Composition**

Ingredients	Gms / Litre
Yeast extract	5.000
Dextrose	20.000
Agar	12.000
Final pH (at 25°C)	6.6±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 18.5 grams in 500 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C and aseptically add reconstituted contents of one vial of Oxytetra Selective Supplement (FD032) or Genta-Oxy Selective Supplement (FD131). Mix well and pour into sterile Petri plates.

Principle And Interpretation

Acidic media are not completely suitable for counting yeasts and moulds in foods since yeast cells, stressed by heat do not tolerate the acidic conditions necessary to inhibit bacterial contamination. Yeast and mould growth is often limited by the presence of acid-tolerant bacterial flora. Therefore it is evident that more active media and different selective agents are needed in order to deal with various kinds of foodstuffs, incubation conditions and types of microorganisms to be studied. Under certain conditions and when testing certain foods like milk and milk products, the use of oxytetracycline alone was not sufficient to obtain reliable yeast and mould counts. In particular, Mossel et al (1) observed that with proteinaceous foods heavily contaminated with gram-negative rods, it is necessary to use both oxytetracycline and gentamicin (FD131) in order to obtain complete inhibition of the contaminants.

OGYE Media were formulated by Mossel et al for the selective isolation and enumeration of yeast and moulds from foods (1, 2). They found that addition of Oxytetra selective supplement to a neutral pH medium increased the recovery / count of yeast and moulds as compared to acidified medium. Psychotrophic yeasts can also be isolated when gentamicin is also incorporated into the medium (3). ISO Committee (4) has recommended OGYE Media with pH 6.6 ± 0.2 for isolation and enumeration of yeasts and moulds from milk and milk products.

Yeast extract provides essential growth nutrients. Dextrose acts as carbon and energy source. Low pH helps to reduce the bacterial flora. Oxytetracycline makes the medium more selective by inhibiting the growth of Lactobacilli encountered in milk and milk-products at low pH. The choice of a suitable media for enumeration of yeasts and moulds greatly depends on the nature of foodstuffs to be tested and the organisms that grow on them. These media remain bacteriostatic when inoculated with not greater than 1 ml of a 10^{-1} food dilution and incubation at 22°C. The number of yeasts or moulds is calculated per one gram or 1 ml of sample under investigation by multiplying the number of colonies with the dilution factor. Lactic acid bacteria are inhibited on this medium.

Quality Control

Appearance

Cream to light yellow coloured granular medium

Gelling

Firm, comparable with 1.2% Agar gel.

Colour and Clarity of prepared medium

Light amber coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Please refer disclaimer Overleaf.

Reaction of 3.7% w/v aqueous solution at 25°C. pH : 6.6±0.2

pH

6.40-6.80

Cultural Response

Cultural characteristics observed with added 1 vial of Oxytetra Selective Supplement(FD032) or Genta-Oxy Selective Supplement (FD131), after an incubation at 25-30°C after 2-5 days.

Cultural Response

Organism	Inoculum (CFU)	Growth	Recovery
Cultural Response			
* <i>Aspergillus brasiliensis</i> ATCC 16404	50-100	good-luxuriant	
<i>Candida albicans</i> ATCC 10231	50-100	good-luxuriant	≥50%
<i>Escherichia coli</i> ATCC 25922	≥10 ³	inhibited	0%
<i>Saccharomyces cerevisiae</i> ATCC 9763	50-100	good-luxuriant	≥50%
<i>Saccharomyces uvarum</i> ATCC 9080	50-100	good-luxuriant	≥50%

*Key: Formerly known as *Aspergillus niger*

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

- 1.Mossel D. A. A., Kleynen-Semmeling H. M., Vincentie H., Beerens H. and Catsaras M., 1970, J. Appl. Bacteriol., 33:454
- 2.Mossel D. A. A., Visser M. and Mengerink W. H. J., 1962, Lab. Pract. 11:109.
- 3.Mossel D. A. A., Vega Clara L. and Put H. M. C., 1975, J. Appl. Bacteriol., 39:15.
- 4.International Organization for Standardization (ISO), 1992, ISO/DIS 6611.

Revision : 00 / 2014



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.