

Iron Sulphite Agar, Granulated

GM868

Iron Sulphite Agar, granulated is recommended for the detection of thermophilic anaerobic organisms causing sulphide spoilage in food.

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	10.000
Sodium sulphite	0.500
Iron (III) citrate	0.500
Agar	15.000
Final pH (at 25°C)	7.1±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 26 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and pour into sterile tubes or Petri plates or as desired.

Principle And Interpretation

Iron Sulphite Agar is a modification of Cameron Sulphite Agar developed by the National Canners Association of America (1). It was shown by Beerens (2) that 0.1% sulphite concentration in the original formula was inhibitory to some strains of *Clostridium sporogenes*. This observation was later confirmed by Mossel et al (3), who consequently showed that 0.05% sulphite concentration was not inhibitory to the organisms. Most clostridia have sulfite reductase in their cytoplasm but they are unable to expel them to the exterior. So when H₂S is produced from sulfite, the colony becomes dark due to the formation of precipitates of iron sulfide from citrate.

Casein enzymic hydrolysate provides nitrogen and other nutrients necessary to support bacterial growth. Sulphite-reducing bacteria usually produce black colonies as a result of the reduction of sulphite to sulphide, which reacts with the iron (III) salt.

For the detection of organisms causing sulphide spoilage, two methods can be followed:

a) Deep-Shake Culture Method: Dispense the medium in 10 ml amounts in tubes. Inoculate the sample when the medium is at about 45-50°C. Allow to set and incubate at 55°C for 24-48 hours.

Typical thermophilic species - *Desulfotomaculum nigrificans*, produces distinct black spherical colonies in the depth of the medium.

b) Attenborough and Scarr (4) Method: In this method, diluted samples of sugar or any other food are filtered through membrane filters. These filters are then rolled up and placed in tubes containing just sufficient Iron Sulphite Agar (at 45-50°C) to cover them. The medium is allowed to set and then incubated at 55-56°C for 24-48 hours. After incubation, the number of black colonies on the membrane filter is counted. Confirmation tests are further carried out to identify the organism growing in the medium. This membrane filter technique is quicker, of comparable accuracy and permits the examination of larger samples. The blackening reaction is only presumptive evidence of clostridial growth. Confirmation test must be carried out for identification. There are many gram-negative bacteria that are able to reduce sulfite with iron sulfide production in this medium, but in these cases the enzymes are extra cellular and the entire medium becomes dark, rendering their enumeration impossible.

Quality Control

Appearance

Light yellow to brownish yellow coloured granular medium

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Please refer disclaimer Overleaf.

Yellow coloured, slightly opalescent gel forms in Petri plates

Reaction

Reaction of 2.6% w/v aqueous solution at 25°C. pH : 7.1±0.2

pH

6.90-7.30

Cultural Response

Cultural characteristics observed under anaerobic conditions, after an incubation at 55-56°C for 24-48 hours.

Cultural Response

Organism	Inoculum	Growth	Recovery	Colour of colony
Cultural Response				
<i>Clostridium botulinum</i> ATCC 25763	50-100	luxuriant	>=50%	black
<i>Clostridium butyricum</i> ATCC 13732	50-100	luxuriant	>=50%	black
<i>Clostridium sporogenes</i> ATCC 19404	50-100	luxuriant	>=50%	black
<i>Desulfotomaculum</i> <i>nigrificans</i> ATCC 19998	50-100	luxuriant	>=50%	black
<i>Escherichia coli</i> ATCC 25922	50-100	good	40-50%	no blackening

Storage and Shelf Life

Store below 30°C in tightly closed container and prepared medium at 2-8°C. Use before expiry period on the label.

Reference

- 1.Tanner F. W., 1944, "The Microbiology of Foods", 2nd Ed., Garrard Press, Illinois, P. 1127.
- 2.Beerens H., 1958, DSIR, Proc. 2nd Internat. Sym. Food Microbiol., 1957, HMSO, London, P. 235.
- 3.Mossel D. A. A., Golstein Brouwers G. W. M. V. and de Bruin A. S., 1959, J. Path. Bacteriol., 78:290.
- 4.Attenborough J. and Scarr M., 1957, J. Appl. Bacteriol., 20: 460.

Revision : 00 / 2014



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.