



Sabouraud Dextrose Agar

M063

Intended Use:

Sabouraud Dextrose Agar is used for the cultivation of yeasts, moulds and aciduric bacteria from clinical and non clinical samples.

Composition**

Ingredients	Gms / Litre
Dextrose (Glucose)	40.000
Mycological, peptone	10.000
Agar	15.000
Final pH (at 25°C)	5.6±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 65.0 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and pour into sterile Petri plates.

Principle And Interpretation

Sabouraud Dextrose Agar is Carlier's modification (1) of the formulation described by is a modification of Sabouraud Dextrose Agar which is described by Sabouraud (2) for the cultivation of fungi (yeasts, moulds), particularly useful for the fungi associated with skin infections. This medium is also employed to determine microbial contamination in food, cosmetics, and clinical specimens (3).

Mycological Peptone provides nitrogenous compounds. Dextrose provides an energy source. High dextrose concentration and low pH favours fungal growth and inhibits contaminating bacteria from test samples (4).

Type of specimen

Clinical samples: skin scrapings , Food samples ; Cosmetics

Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (7,8).

For food and dairy samples, follow appropriate techniques for sample collection and processing as per guidelines(5,6,9). After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions:

In Vitro diagnostic use only. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations

1. For heavily contaminated samples, the plate must be supplemented with inhibitory agents for inhibiting bacterial growth with lower pH.
2. Some pathogenic fungi may produce infective spores which are easily dispersed in air, so examination should be carried out in safety cabinet
3. Further biochemical tests should be carried out for confirmation.

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder.

Gelling

Firm, comparable with 1.5% Agar gel.

Colour and Clarity of prepared medium

Light amber coloured clear to slightly opalescent gel forms in Petri plates.

Reaction

Reaction of 6.5% w/v aqueous solution at 25°C (after sterilization). pH : 5.6±0.2

pH

5.40-5.80

Cultural Response

Growth Promotion was carried out in accordance with the (USP/EP/BP/JP), after an incubation at 20-25 °C for 24-48 hours. Recovery rate is considered as 100% for bacteria growth on Soybean Casein Digest Agar and fungus growth on Sabouraud Dextrose Agar

Growth Promotion Test

Growth Promotion was carried out in accordance with the harmonized method of ICH (USP/EP/BP/JP), after an incubation at 30-35 °C for 24-48 hours. Recovery rate is considered as 100% for bacteria growth on Soybean Casein Digest Agar and fungus growth on Sabouraud Dextrose Agar

Growth Promoting Properties

Growth of microorganism comparable to that previously obtained with previously tested and approved lot of medium occurs at the specified temperature for not more than the shortest period of time specified inoculating ≥ 100 cfu (at 30-35°C for 24 hours).

Indicative properties

Colonies are comparable in appearance and indication reaction to those previously obtained with previously tested and approved lot of medium occurs for the specified temperature for a period of time within the range specified inoculating ≥ 100 cfu (at 30-35°C for 24-48 hours).

Cultural Response

Organism	Inoculum (CFU)	Growth	Recovery
<i>Candida albicans</i> ATCC 10231 (00054*)	50 -100	Luxuriant (white colonies)	≥ 70 %
<i>Aspergillus brasiliensis</i> ATCC 16404 (00053*)	50 -100	luxuriant	≥ 70 %
<i>Candida albicans</i> ATCC 2091 (00055*)	50 -100	luxuriant	≥ 70 %
<i>Saccharomyces cerevisiae</i> ATCC 9763 (00058*)	50 -100	luxuriant	≥ 70 %
<i>Escherichia coli</i> ATCC 8739 (00012*)	50 -100	luxuriant	≥ 70 %
<i>Escherichia coli</i> ATCC 25922 (00013*)	50 -100	luxuriant	≥ 70 %
<i>Escherichia coli</i> NCTC 9002	50 -100	luxuriant	≥ 70 %
<i>Lactobacillus casei</i> ATCC 334	50 -100	luxuriant	≥ 70 %
<i>Trichophyton rubrum</i> ATCC 28191		luxuriant	

Key : *Corresponding WDCM numbers.

Storage and Shelf Life

Store below 30°C in a tightly closed container and the prepared medium at 20-30°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Use before expiry date on the label. Product performance is best if used within stated expiry period.

Please refer disclaimer Overleaf.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (7,8).

Reference

1. Carlier G. I. M., 1948, Brit. J. Derm. Syph., 60:61.
2. Sabouraud K., 1892, Ann. Dermatol. Syphilol, 3:1061.
3. Bacteriological Analytical Manual, 8th Edition, Revision A, 1998. AOAC, Washington D.C.
4. Murray PR, Baren EJ, Jorgensen JH, Pfaller MA, Tenover FC, Tenover FC (editors) 2003, Manual of clinical Microbiology, 8th ed.,ASM, Washington, D.C.
5. American Public Health Association, Standard Methods for the Examination of Dairy Products, 1978, 14th Ed., Washington D.C.
6. Salfinger Y., and Tortorello M.L. Fifth (Ed.), 2015, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.
7. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
8. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock, D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
9. Wehr H. M. and Frank J. H., 2004, Standard Methods for the Microbiological Examination of Dairy Products, 17th Ed., APHA Inc., Washington, D.C.


Revision : 03 / 2018



In vitro diagnostic medical device



CE Marking



Storage temperature



Do not use if package is damaged



HiMedia Laboratories Pvt. Limited,
23 Vadhani Industrial Estate,
LBS Marg, Mumbai-86, MS, India



CE Partner 4U, Esdoornlaan 13, 3951
DB Maarn The Netherlands,
www.cepartner4u.eu

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.