



Corn Meal Agar

M146

Corn Meal Agar is recommended for chlamyospore production by *Candida albicans* and the maintenance of fungal stock cultures.

Composition**

Ingredients	Gms / Litre
Corn meal, infusion from	50.000
Agar	15.000
Final pH (at 25°C)	6.0±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 17 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. If desired add 1% polysorbate 80. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour into sterile Petri plates.

Principle And Interpretation

Chlamyospore production is an accepted criterion for the identification of *Candida* species. Corn Meal Agar is a well-established mycological medium used for the cultivation of fungi and to study chlamyospores production of *Candida* species (1). Corn Meal Agar is a general purpose medium used for the cultivation of fungi and for the study of *Candida* species for chlamyospore production. Pollack and Benham (1) have described the usefulness of this medium for studying the morphology of *Candida*. Walker and Huppert (2) modified this medium by adding polysorbate 80, which then stimulated faster and plenty of chlamyospore formation of *Candida* species.

This is a very simple formulation containing only cornmeal infusion and agar. However this infusion has enough nutrients to enhance the growth of fungi. Polysorbate 80 is a mixture of oleic esters, which activates the production of chlamyospore by *Candida albicans*, *Candida stellatoidea* and *Candida tropicalis* (3). Some *Candida* species lose their ability of chlamyospore formation by repeated sub culturing.

Pick a suspected colony from Sabouraud Dextrose Agar (M063) using a straight wire, and make a deep cut in the Corn Meal Agar plate. Repeat for each colony. Place a flamed sterile coverslip over the line of inoculum. After incubation for 24-48 hours at 25-30°C, the streaks are examined microscopically, through the coverslip, using low and high power objectives.

C. albicans produces mycelium bearing ball-like clusters of budding cells and characteristics thick walled round chlamyospores (4).

Quality Control

Appearance

Cream to yellow coarse free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Light amber coloured, opalescent gel forms in Petri plates

Reaction

Reaction of 1.7% w/v aqueous solution at 25°C. pH : 6.0±0.2

pH

5.80-6.20

Cultural Response

M146: Cultural characteristics observed after an incubation at 23-27°C for upto 4 days.

Organism	Inoculum (CFU)	Growth	Chlamyospores	Recovery
----------	-------------------	--------	---------------	----------

* <i>Aspergillus brasiliensis</i> ATCC 16404	50-100	luxuriant	negative	
<i>Candida albicans</i> ATCC 10231	50-100	luxuriant	positive	>=70%
<i>Saccharomyces cerevisiae</i> ATCC 9763	50-100	luxuriant	negative	>=70%
<i>Saccharomyces uvarum</i> ATCC 28098	50-100	luxuriant	negative	>=70%

Key : * - Formerly known as *Aspergillus niger*

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

Reference

1. Pollack and Benham, 1960, J. Lab. Clin. Med., 50:313.
2. Walker and Huppert, 1960, Tech. Bull. Reg. Med. Technol., 30:10.
3. Cooper and Silvo-Hunter, 1985, Manual of Clinical Microbiology, Lennette, Balows, Hausler and Shadomy (Eds.), 4th ed., ASM, Washington, D.C.
4. Conant N. F., Smith D. T., Baker R. D., Callaway J. L. and Martin D. S., 1971, Manual of Clinical Mycology, 3rd Ed., USA

Revision : 2 / 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.