



## Actidione Agar Base w/o Actidione®

M058

### Intended Use:

Recommended for the enumeration and detection of bacteria in specimens containing large numbers of yeasts and moulds.

### Composition\*\*

Ingredients	Gms / Litre
Tryptone	5.000
Yeast extract	4.000
Dextrose (Glucose)	50.000
Potassium dihydrogen phosphate	0.550
Potassium chloride	0.425
Calcium chloride anhydrous	0.125
Magnesium sulphate	0.125
Ferric chloride	0.0025
Manganese sulphate	0.0025
Bromo cresol green	0.022
Agar	15.000
Final pH ( at 25°C)	5.5±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 75.25 grams in 1000 ml purified / distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C and aseptically add 10 mg to the medium. Mix well before pouring into sterile Petri plates.

Warning : Cycloheximide (Actidione) is very toxic. Avoid skin contact or aerosol formation and inhalation.

### Principle And Interpretation

Actidione Agar was formulated by Green and Gray (1) which may be used for microbiological investigation during brewing and baking. Actidione® (Cycloheximide) at a concentration of 0.001% permits the growth of bacteria and inhibits the growth of most yeasts and moulds except dermatophytes. This medium may be used for the estimation of bacterial contamination of pitching yeast. Addition of penicillin or streptomycin may be used for selective isolation of dermatophytes.

Tryptone acts as source of nitrogen while yeast extract is a rich reservoir of vitamins. Dextrose in high amount along with mineral salts at acidic pH favours sugar fermentation.

### Type of specimen

Brewing sample.

### Specimen Collection and Handling

For Brewing samples, follow appropriate techniques for sample collection, processing as per guidelines and local standards. (1).

After use, contaminated materials must be sterilized by autoclaving before discarding.

### Warning and Precautions

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

### Limitations

1. Further Biochemical tests must be carried out for confirmation.
2. Cycloheximide (Actidione) is very toxic. Avoid skin contact or aerosol formation and inhalation.

## Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

## Quality Control

### Appearance

Light yellow to light green homogeneous free flowing powder

### Gelling

Firm, comparable with 1.5% Agar gel

### Colour and Clarity of prepared medium

Greenish blue clear to slightly opalescent gel forms in Petri plates

### Reaction

Reaction of 7.53% w/v aqueous solution at 25°C. pH : 5.5±0.2

### pH

5.30-5.70

### Cultural Response

Cultural characteristics observed after an incubation at 25-30°C for 40-48 hours.

Organism	Inoculum (CFU)	Growth (Plain)	Recovery	Growth (w/ Actidione)	Recovery (w / Actidione)
<i>Escherichia coli</i> ATCC 25922	50-100	luxuriant	≥70%	good-luxuriant	≥50%
<i>Lactobacillus fermentum</i> ATCC 9338	50-100	luxuriant	≥70%	good-luxuriant	≥50%
<i>Proteus mirabilis</i> ATCC 25933	50-100	luxuriant	≥70%	good-luxuriant	≥50%
<i>Saccharomyces cerevisiae</i> ATCC 9763	50-100	luxuriant	≥70%	inhibited	0%
<i>Saccharomyces uvarum</i> ATCC 9080	50-100	luxuriant	≥70%	inhibited	0%

Key : (\*) Corresponding WDCM numbers.

## Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Use before expiry date on the label.

Product performance is best if used within stated expiry period.

## Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (2,3).

## Reference

- Green, S.R. and Gray, P.P., 1950, Wallerstein Lab. Communication, 13:357.
- Isenberg, H.D. Clinical Microbiology Procedures Handbook 2<sup>nd</sup> Edition.
- Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision :02 / 2019

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.