

## Medium 199 (For Suspension Culture)

With Earle's salts and L-Glutamine

Without Calcium chloride, Magnesium sulphate and Sodium bicarbonate

**Product Code: AT043**

### Product Description :

Medium 199 was the first nutritionally defined medium developed by Morgan, Morton, and Parker in 1950. This complex medium was formulated specifically for nutritional studies on primary chick embryo fibroblasts in the absence of any additives. It was observed that explanted tissue could survive in Medium 199 without serum but long term cultivation of cells required supplementation of the medium with serum.

Medium 199 is formulated with either Hank's salts or Earle's salts. The medium when supplemented with serum can be used for growth of a wide variety of cells. Medium 199 is presently used for the maintenance of non-transformed cells, vaccine and virus production and primary explants of epithelial cells.

AT043 is Medium 199 with Earle's salts and L-Glutamine. It is modified for use with suspension cells, hence does not contain calcium chloride and magnesium sulphate. Users are advised to review the literature for recommendations regarding medium supplementation and physiological growth requirements specific for different cell lines.

### Composition :

Ingredients	mg/L
<b>INORGANIC SALTS</b>	
Ferric nitrate nonahydrate	0.720
Potassium chloride	400.000
Sodium acetate anhydrous	50.000
Sodium chloride	6800.000
Sodium phosphate monobasic	122.000
<b>AMINO ACIDS</b>	
Glycine	50.000
L-Alanine	25.000
L-Arginine hydrochloride	70.000
L-Aspartic acid	30.000
L-Cysteine hydrochloride monohydrate	0.100
L-Cystine dihydrochloride	26.000
L-Glutamic acid	67.000
L-Glutamine	100.000

L-Histidine hydrochloride monohydrate	22.000
L-Hydroxyproline	10.000
L-Isoleucine	20.000
L-Leucine	60.000
L-Lysine hydrochloride	70.000
L-Methionine	15.000
L-Phenylalanine	25.000
L-Proline	40.000
L-Serine	25.000
L-Threonine	30.000
L-Tryptophan	10.000
L-Tyrosine disodium salt	57.660
L-Valine	25.000
<b>VITAMINS</b>	
Ascorbic acid	0.050
Calciferol	0.100
Choline chloride	0.500
D-Biotin	0.010
D-Ca-Pantothenate	0.010
DL-Tocopherol phosphate disodium salt	0.010
Folic acid	0.010
Menadione	0.010
Nicotinamide	0.025
Nicotinic acid	0.025
Pyridoxal hydrochloride	0.025
Pyridoxine hydrochloride	0.025
Retinol Acetate	0.140
Riboflavin	0.010
Thiamine hydrochloride	0.010
i-Inositol	0.050
p-Amino benzoic acid (PABA)	0.050
<b>OTHERS</b>	
Adenine sulphate	10.000
Adenosine monophosphate	0.200
Adenosine triphosphate	1.000
Cholesterol	0.200
Deoxyribose	0.500
Glucose	1000.000
Glutathione reduced	0.050
Guanine hydrochloride	0.300

Hypoxanthine	0.354
Phenol red	15.000
Polysorbate 80	4.900
Ribose	0.500
Thymine	0.300
Uracil	0.300
Xanthine	0.344

### Directions :

1. Suspend 9.2gms in 900ml tissue culture grade water with constant, gentle stirring until the powder is completely dissolved. Do not heat the water.
2. Add 2.2gms of sodium bicarbonate powder (TC230) or 29.3ml of 7.5% sodium bicarbonate solution (TCL013) for 1 litre of medium and stir until dissolved.
3. Adjust the pH to 0.2 - 0.3 pH units below the desired pH using 1N HCl or 1N NaOH since the pH tends to rise during filtration.
4. Make up the final volume to 1000ml with tissue culture grade water.
5. Sterilize the medium immediately by filtering through a sterile membrane filter with a porosity of 0.22 micron or less, using positive pressure rather than vacuum to minimize the loss of carbon dioxide.
6. Aseptically add sterile supplements as required and dispense the desired amount of sterile medium into sterile containers.
7. Store liquid medium at 2-8°C and in dark till use.

### Material required but not provided :

Tissue culture grade water (TCL010)  
Sodium bicarbonate (TC230)  
Sodium bicarbonate solution, 7.5% (TCL013)  
1N Hydrochloric acid (TCL003)  
1N Sodium hydroxide (TCL002)  
Foetal bovine serum (RM1112/RM10432)

### Quality Control:

#### Appearance

Off-white to Creamish white, homogenous powder.

#### Solubility

Clear solution at 9.2 gms/L.

#### pH without Sodium Bicarbonate

4.30 -5.00

#### pH with Sodium Bicarbonate

7.10 -7.70

### Osmolality without Sodium Bicarbonate

230.00 -270.00

### Osmolality with Sodium Bicarbonate

280.00 -320.00

### Cultural Response

The growth promotion capacity of the medium is assessed qualitatively by analyzing the cells for the morphology and quantitatively by estimating the cell counts and comparing it with a control medium through minimum three subcultures.

### Endotoxin content

NMT 5EU/ml

### Storage and Shelf Life:

1. All the powdered media and prepared liquid culture media should be stored at 2-8°C. Use before the expiry date. In spite of above recommended storage condition, certain powdered medium may show some signs of deterioration /degradation in certain instances. This can be indicated by change in colour, change in appearance and presence of particulate matter and haziness after dissolution.
2. Preparation of concentrated medium is not recommended since free base amino acids and salt complexes having low solubility may precipitate in concentrated medium.
3. pH and sodium bicarbonate concentration of the prepared medium are critical factors affecting cell growth. This is also influenced by amount of medium and volume of culture vessel used (surface to volume ratio). For example, in large bottles, such as Roux bottles pH tends to rise perceptibly as significant volume of carbon dioxide is released. Therefore, optimal conditions of pH, sodium bicarbonate concentration, surface to volume ratio must be determined for each cell type. We recommend stringent monitoring of pH. If needed, pH can be adjusted by using sterile 1N HCl or 1N NaOH or by bubbling in carbon dioxide.
4. If required, supplements can be added to the medium prior to or after filter sterilization observing sterility precautions. Shelf life of the medium will depend on the nature of supplement added to the medium.

Revision : 1 / 2012

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

