



# Technical Data

## HiEncap™ Sabouraud Dextrose Broth (HiEncap™ Sabouraud EC033CCL Liquid Medium)

HiEncap™ Sabouraud Dextrose Broth (Sabouraud Liquid Medium) is used for cultivation of yeasts, moulds and aciduric microorganisms.

### Composition\*\*

Ingredients	Gms / Litre
Dextrose	20.000
Peptone, special	10.000
Final pH ( at 25°C)	5.6±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Each capsule contains 7.5 grams of media. Suspend 1 capsule in 250 ml (4 capsules in 1000 ml) distilled or purified water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and dispense as desired.

### Principle And Interpretation

Sabouraud Dextrose Agar is Carliers modifications (1) of the formulation described by Sabouraud (2) for the cultivation of fungi, particularly those associated with skin infections. The medium is also recommended by APHA (3). Sabouraud Dextrose Broth is also a modification by Sabouraud (4) and serves the same purpose as Sabouraud Dextrose Agar.

Sabouraud dextrose media are peptone media supplemented with dextrose to support the growth of fungi. Peptone special provides nitrogen, vitamins, minerals, amino acids and growth factors. Dextrose provides an energy source for the growth of microorganisms. The low pH favours fungal growth and inhibits contaminating bacteria from clinical specimens (5). The acid reaction of the final medium is inhibitory to a large number of bacteria making it particularly useful for cultivating fungi and aciduric microorganisms. For isolation of fungi from contaminated specimens, a selective medium should be inoculated simultaneously. Incubate cultures for 4 to 6 weeks before reporting as negative.

### Quality Control

#### Appearance

Gelatin capsule containing cream to yellow coloured granular media

#### Colour and Clarity of prepared medium

Light amber coloured clear solution in tubes

#### Quantity

Each capsule contains 7.5 grams of medium sufficient for 250 ml media

#### Reaction

pH of 3.0% w/v aqueous solution at 25°C. pH : 5.6±0.2

#### pH

5.40-5.80

#### Cultural Response

Cultural characteristics was observed after an incubation at 20-25°C for 3-5 days.

#### Cultural Response

EC033CCL: Cultural characteristics observed after incubation at 20-25 °C for 3-5 days.

Organism	Inoculum (CFU)	Growth	Incubation temperature	Incubation period
<b>Cultural Response</b> <i>Candida albicans</i> ATCC 10231	50 -100	luxuriant	20 -25 °C	<=5 d

<i>*Aspergillus brasiliensis</i> ATCC 16404	50 -100	luxuriant	20 -25 °C	<=5 d
<i>Saccharomyces cerevisiae</i> ATCC 9763	50 -100	luxuriant	20 -25 °C	3 -5 d
<i>Saccharomyces cerevisiae</i> ATCC 2601	50 -100	good-luxuriant	20 -25 °C	3 -5 d
<i>Candida albicans</i> ATCC 2091	50 -100	luxuriant	20 -25 °C	3 -5 d
<i>Escherichia coli</i> ATCC 8739	50 -100	Luxuriant (inhibited on media with low pH)	20 -25 °C	<=5 d
<i>Escherichia coli</i> ATCC 25922	50 -100	good-luxuriant	20 -25 °C	3 -5 d
<i>Escherichia coli</i> NCTC 9002	50 -100	Luxuriant (inhibited on media with low pH)	20 -25 °C	3 -5 d
<i>Lactobacillus casei</i> ATCC 334	50 -100	luxuriant	20 -25 °C	3 -5 d

### Storage and Shelf Life

Store below 30°C in a tightly closed container and prepared medium at 2 - 8°C. Use before expiry date on the label.

### Reference

1. Carlier G. I. M., 1984, Brit. J. Derm. Syph., 60:61
2. Sabouraud R., 1892, Ann. Dermatol. Syphil. 3 : 1061.
3. Downes F. P. and Ito K., (Eds.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., APHA, Washington, D.C.
4. Sabouraud R., Les Teignes, Paris: Masson et Cie, 1910, p 553
5. Murray P. R., Baron J. H., Pfaller M. A., Jorgensen J. H. and Tenover F. C., (Ed.), 2003, Manual of Clinical Microbiology, 8th Ed., American Society for Microbiology, Washington, D.C.

Revision : 00 / 2014



#### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.