



# Technical Data

## HiEncap™ Plate Count Agar (HiEncap™ Standard Methods EC091CCL Agar)

HiEncap™ Plate Count Agar is recommended for the determination of plate counts of microorganisms in food, water, waste water and clinical samples.

### Composition\*\*

Ingredients	Gms / Litre
Casein enzymic hydrolysate	5.000
Yeast extract	2.500
Dextrose	1.000
Agar	15.000
Final pH ( at 25°C)	7.0±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Each capsule contains 5.88 gms of medium. Suspend 1 capsule in 250 ml (4 capsules in 1000 ml) distilled or purified water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour into sterile Petri plates.

### Principle And Interpretation

Plate Count Agar is formulated as described by Buchbinder et al (1) which is recommended by APHA (2,3,4) and FDA (5).

Casein enzymic hydrolysate provides amino acids and other complex nitrogenous substances. Yeast extract supplies Vitamin B complex. APHA recommends the use of pour plate technique. The samples are diluted and appropriate dilutions are added in Petri plates. Sterile molten agar is added to these plates and plates are rotated gently to ensure uniform mixing of the sample with agar. The poured plate count method is preferred to the surface inoculation method, since it gives higher results. Plate Count Agar is also suitable for enumerating bacterial count of sterile rooms.

### Quality Control

#### Appearance

Gelatin capsule containing cream to yellow coloured granulated media.

#### Gelling

Firm, comparable with 1.5% Agar gel

#### Colour and Clarity of prepared medium

Light yellow coloured clear to slightly opalescent gel forms in Petri plates

#### Quantity

Each capsule contains 5.88 grams of medium sufficient for 250 ml media

#### Reaction

Reaction of 2.35% w/v aqueous solution at 25°C. pH : 7.0±0.2

#### pH

6.80-7.20

#### Cultural Response

Cultural characteristics observed after an incubation at 35 - 37°C for 18 - 48 hours.

#### Cultural Response

Organism	Growth	Inoculum (CFU)	Recovery
----------	--------	----------------	----------

#### Cultural Response

<i>Lactobacillus casei</i> ATCC 9595	luxuriant	50-100	>=70%
--------------------------------------	-----------	--------	-------

---

<i>Staphylococcus aureus</i> ATCC 25923	luxuriant	50-100	>=70%
<i>Streptococcus pyogenes</i> ATCC 19615	luxuriant	50-100	>=70%
<i>Bacillus subtilis</i> ATCC 6633	luxuriant	50-100	>=70%
<i>Enterococcus faecalis</i> ATCC 29212	luxuriant	50-100	>=70%
<i>Escherichia coli</i> ATCC 25922	luxuriant	50-100	>=70%

## Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

## Reference

1. Buchbinder L., Baris Y., Aldd E., Reynolds E., Dilon E., Pessin V., Pincas L. and Strauss A., 1951, Publ. Hlth. Rep., 66:327.
2. Wehr H. M. and Frank J. H., 2004, Standard Methods for the Microbiological Examination of Dairy Products, 17th Ed., APHA Inc., Washington, D.C.
3. Downes F. P. and Ito K., (Eds.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., APHA, Washington, D.C.
4. Eaton A. D., Clesceri L. S. and Greenberg A. E., Rice E. W., (Eds.), 2005, Standard Methods for the Examination of Water and Wastewater, 21st Ed., APHA, Washington, D.C.
5. FDA Bacteriological Analytical Manual, 2005, 18th Ed., AOAC, Washington, DC.

Revision : 00 / 2014



### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.