



# Technical Data

## Potassium Tellurite 3.5% (1 ml per vial)

FD047

Recommended for the selective isolation of Staphylococci and Corynebacteria.

### Composition

Per vial sufficient for medium as specified in direction.

#### Ingredients

Potassium tellurite  
Distilled water

#### Concentration

0.350g  
1ml

### Directions:

Warm up the refrigerated contents of one vial to 45-50°C. Add aseptically 9 ml sterile distilled water, mix well and add 3 ml in 950 ml sterile, molten, cooled (45-50°C) Baird Parker Agar Base [M043](#) / [M043S](#) / Baird Parker HiVeg™ Agar Base [MV043](#) / Baird Parker Agar Base w/ Sulpha [M1140](#) along with 50 ml Concentrated Egg Yolk Emulsion [FD045](#) or 10 ml in 1000 ml Hoyle Medium Base [M015](#) / Hoyle HiVeg™ Medium Base [MV015](#) alongwith 50 ml of laked blood. Mix well and pour into sterile petri plates.

### Storage and Shelf Life

Store at 2-8°C. Use before expiry period on the label.

#### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.