

## Phenol Red Broth Base, Granulated

GM054

Phenol Red Broth Base, granulated is used as a basal medium to which carbohydrates are added for determination of fermentation reactions of pure cultures of microorganisms.

### Composition\*\*

Ingredients	Gms / Litre
Proteose peptone	10.000
Beef extract	1.000
Sodium chloride	5.000
Phenol red	0.018
Final pH ( at 25°C)	7.4±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 16.02 grams in 1000 ml purified / distilled water. Heat if necessary to to dissolve the medium completely. Mix well and dispense in tubes containing inverted Durham's tubes and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Aseptically add filter sterilized or autoclave sterilized desired amount of carbohydrate solution to sterile basal medium.

### Principle And Interpretation

Phenol Red Broth Medium is formulated as per Vera (2) and is recommended to determine the fermentation reaction of carbohydrates for the differentiation of microorganisms (3, 4, 5). Phenol Red Broth Medium with various added carbohydrates serves as a differential medium by aiding in differentiation of various species and genera by their ability to ferment the specific carbohydrate, with the production of acid or acid and gas (6).

Phenol Red Broth Base is a complete medium without added carbohydrate, which can be used with the addition of 5-10 %, desired carbohydrate. It is used as a negative control for studying fermentations or as a base for the addition of carbohydrates. Proteose peptone and beef extract serve as sources for carbon and nitrogen. Sodium chloride is the osmotic stabilizer. Phenol red is the pH indicator, which turns yellow at acidic pH. Gas formation is seen in Durhams tubes. All of the *Enterobacteriaceae* grow well in this medium. In addition to producing a pH colour shift, the production of mixed acids, notably butyric acids, often results in a pungent, foul odour from the culture medium (1).

### Quality Control

#### Appearance

Light yellow to pink coloured granular medium

#### Colour and Clarity of prepared medium

Red coloured clear solution without any precipitate

#### Reaction

Reaction of 1.6% w/v aqueous solution at 25°C. pH : 7.4±0.2

#### pH

7.20-7.60

#### Cultural Response

Cultural characteristics observed after an incubation at 35 - 37°C for 18 - 24 hours.( longer if necessary)

Organism	Inoculum (CFU)	Growth	without carbohydrate, (Acid)	without carbohydrate, (Gas)	with dextrose, (Acid)	with dextrose, (Gas)
<b>Cultural Response</b>						
<i>Citrobacter freundii</i> ATCC 8090	50-100	luxuriant	Negative reaction, no colour change	Negative reaction	Positive reaction, yellow colour	Positive reaction

<i>Escherichia coli</i> ATCC 25922	50-100	luxuriant	Negative reaction, no colour change	Negative reaction	Positive reaction, yellow colour	Positive reaction
<i>Enterobacter aerogenes</i> ATCC 13048	50-100	luxuriant	Negative reaction, no colour change	Negative reaction	Positive reaction, yellow colour	Positive reaction
<i>Klebsiella pneumoniae</i> ATCC 13883	50-100	luxuriant	Negative reaction, no colour change	Negative reaction	Positive reaction, yellow colour	Positive reaction
<i>Proteus vulgaris</i> ATCC 13315	50-100	luxuriant	Negative reaction, no colour change	Negative reaction	Positive reaction, yellow colour	Positive reaction
<i>Salmonella</i> Typhi ATCC 6539	50-100	luxuriant	Negative reaction, no colour change	Negative reaction	Positive reaction, yellow colour	Positive reaction
<i>Salmonella</i> Typhimurium ATCC 14028	50-100	luxuriant	Negative reaction, no colour change	Negative reaction	Positive reaction, yellow colour	Positive reaction
<i>Serratia marcescens</i> ATCC 8100	50-100	luxuriant	Negative reaction, no colour change	Negative reaction	Positive reaction, yellow colour	Positive reaction
<i>Shigella flexneri</i> ATCC 12022	50-100	luxuriant	Negative reaction, no colour change	Negative reaction	Positive reaction, yellow colour	Negative reaction

## Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

## Reference

1. Koneman E. W., Allen S. D., Janda W.M., Schreckenberger P.C., Winn W.C. Jr., 1992, Colour Atlas and Textbook of Diagnostic Microbiology, 4th Ed., J. B. Lippincott Company
2. Vera H. D., 1950, Am. J. Public Health, 40, 1267
3. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification -Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.
4. Finegold S. M. and Baron E. J., 1986, Bailey and Scotts Diagnostic Microbiology, 7th Ed., The C.V. Mosby Co., St. Louis.
5. Ewing W. H., 1986, Edwards and Ewings Identification of Enterobacteriaceae, 4th ed., Elsevier Science Publishing Co., Inc., New York.
6. MacFaddin J. F., 2000, Biochemical tests for Identification of Medical Bacteria, 3rd ed., Lippincott, Williams and Wilkins, Baltimore.

Revision : 01 / 2014



### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.