

Rinsing Fluid, Granulated

GM1622

This medium is used as a rinsing fluid in the membrane filtration procedure.

Composition**

Ingredients	Gms / Litre
Peptone from meat	5.000
Meat extract	3.000
Polysorbate 80	1.000
Final pH (at 25°C)	6.9±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 9 grams in 1000 ml distilled water. If desired add additional 9 gram quantity of Polysorbate 80 (Tween® 80), filter until clear. Mix well and dispense in tubes or flasks as desired. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Principle And Interpretation

After filtration it is often necessary to rinse the membrane filter in order to remove residues of liquid sample materials. If the sample contains higher hydrocarbons such as vaseline, paraffin, etc. or fats, the use of rinse fluid is recommended. Rinsing fluid largely complies with the formulation prescribed in the recommendations of the United States Pharmacopoeia (1).

This fluid contains balanced concentrations of nutrients, which prevent the microorganisms, retained by the filter, from being exposed to physiological shock, thus being capable to grow further rapidly. The detergent polysorbate 80 ensures emulsification of carbohydrates and fats without seriously affecting the microorganisms. If the sample contains large quantities of these compounds, additional up to 9.0 g/litre of polysorbate 80 (Tween® 80) can be added in accordance with the USP recommendations before the broth is filtered. After filtering the liquid sample, rinse the filter 3 times with 100 ml portions of the membrane-filter rinse fluid, and then complete the test in the usual way.

Quality Control

Appearance

Cream to yellow coloured granular medium

Colour and Clarity of prepared medium

Yellow coloured clear solution without any precipitate

Reaction

Reaction of 0.9% w/v aqueous solution at 25°C pH : 6.9±0.2

pH

6.70-7.10

Cultural Response

Cultural characteristics observed after an incubation at 35 -37°C for 18 - 24 hours.Recovery is carried out on

Standard Nutrient Agar No.1 (GM1210 / M1210)

Organism

Growth

Cultural Response

Staphylococcus aureus Good

ATCC 25923

Streptococcus pyogenes Good

ATCC 12344

Enterococcus faecalis ATCC Good

29212

Citrobacter freundii ATCC Good

8090

Pseudomonas aeruginosa Good

ATCC 27853

Escherichia coli ATCC 25922 Good

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

Reference

1. The United States Pharmacopoeia, 2014, USP37/NF32. The United States Pharmacopoeial Convention. Rockville, MD.

Revision : 00 / 2014



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.