



# Technical Data

## Buffered Sodium Chloride Peptone Solution pH 7.00

LQ123L

For the preparation of test suspension in accordance with harmonized methods of USP, EP, BP, JP & IP

### Composition\*\*

Ingredients	Gms / Litre
Potassium dihydrogen phosphate	3.600
Disodium hydrogen phosphate dihydrate	7.200
Sodium chloride	4.300
HMC peptone #	1.000

\*\*Formula adjusted, standardized to suit performance parameters

# Equivalent to Peptone (meat or casein)

### Directions

Label the ready to use LQ123L bottle. Inoculate the sample and Incubate at specified temperature and time.

### Principle And Interpretation

The composition of this medium is in accordance with the harmonized methodology of USP/EP/BP/JP/IP (7,2,1,5,3). This medium is recommended for preparation of stable test strain suspension employed for validating the microbiological testing procedures of non-sterile products. The standardized stable suspensions are used so that the suitability of this test to detect microorganism in presence of product can be established. Non-fatty products insoluble in water and water-soluble products are diluted/dissolved using this solution.

HMC Peptone serves as nutrient source and maintains the cell viability. Phosphates in the medium act as good buffering agents. Sodium chloride maintains the osmotic balance and cell integrity. Polysorbates reduce surface tension and also inactivate phenolic compound, if present in the test sample.

Preparation of test strain is recommended in Buffered Sodium chloride-Peptone solution pH 7.0 at 30-35°C wherein there is no multiplication of organisms or there is no decrease in count for upto 4 hours. The medium also supports the repair of injured cells that have sensitivity to low pH.

### Type of specimen

Pharmaceutical samples

### Specimen Collection and Handling

For pharmaceutical samples follow appropriate techniques for handling specimens as per established guidelines (7,2,1,5,3).

After use, contaminated materials must be sterilized by autoclaving before discarding.

### Warning and Precautions

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets

### Limitations

This medium contains less nutrients and is not recommended for the growth of microorganisms

### Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

## Quality Control

### Appearance

Sterile clear Buffered Sodium Chloride Peptone Solution in bottle.

### Colour

Colourless solution.

### Quantity of medium

300 ml of medium in bottle

### pH

7.00- 7.00

### Growth Promotion Test

In accordance with the harmonized method of USP/EP/BP/JP.

### Sterility test

Passes release criteria

### Growth Promotion Test

Growth Promotion is carried out in accordance with the harmonized method of (USP/EP/BP/JP/IP).

### Cultural response

Cultural characteristics observed after recovery on Soybean Casein Digest Agar after an incubation at 30-35°C for 18-24 hours for bacteria and Sabouraud Dextrose Agar at 30-35°C for 24-48 hours .

Organism	Inoculum (CFU)	Recovery within 2 hours of incubation	Recovery within 4 hours of incubation	Recovery within 24 hours of incubation
<i>Escherichia coli</i> ATCC 8739 (00012*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Escherichia coli</i> ATCC 25922 (00013*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Escherichia coli</i> NCTC 9002	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 6538 (00032*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 25923 (00034*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Pseudomonas aeruginosa</i> ATCC 9027 (00026*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Pseudomonas aeruginosa</i> ATCC 27853 (00025*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Salmonella Typhimurium</i> ATCC 14028 (00031*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)

<i>Salmonella Abony</i> NCTC 6017 (00029*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Bacillus subtilis</i> subsp. <i>spizizenii</i> ATCC 6633 (00003*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Micrococcus luteus</i> ATCC 9341	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Candida albicans</i> ATCC 10231 (00054*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)
<i>Candida albicans</i> ATCC 2091 (00055*)	50 -100	no decrease in colony count	no decrease in colony count	no decrease in colony count (stored at 2-8°C)

Key : (\*) Corresponding WDCM Numbers

## Storage and Shelf Life

Store between 15-25°C. Use before expiry date on the label.

## Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (4,6).

## Reference

1. British Pharmacopoeia, 2016 The Stationery office British Pharmacopoeia
2. European Pharmacopoeia, 2017, European Dept. for the quality of Medicines.
3. Indian Pharmacopoeia, 2018, Govt. of India, the controller of Publication, Delhi, India.
4. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2<sup>nd</sup> Edition.
5. Japanese Pharmacopoeia, 2016.
6. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
7. The United States Pharmacopoeia, 2019, The United States Pharmacopoeial Convention. Rockville, MD.

Revision : 00 / 2019

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.