



## Xylose-Lysine Deoxycholate Agar (XLD Agar)

M031

### Intended use

Xylose-Lysine Deoxycholate Agar (XLD Agar) is a selective medium recommended for the isolation and enumeration of *Salmonella* Typhi and other *Salmonella* species.

### Composition\*\*

Ingredients	Gms / Litre
Yeast extract	3.000
L-Lysine	5.000
Lactose	7.500
Sucrose	7.500
Xylose	3.500
Sodium chloride	5.000
Sodium deoxycholate	2.500
Sodium thiosulphate	6.800
Ferric ammonium citrate	0.800
Phenol red	0.080
Agar	15.000
Final pH ( at 25°C)	7.4±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 56.68 grams in 1000 ml distilled water. Heat with frequent agitation until the medium boils. DO NOT AUTOCLAVE OR OVERHEAT. Transfer immediately to a water bath at 50°C. After cooling, pour into sterile Petri plates. It is advisable not to prepare large volumes that will require prolonged heating, thereby producing precipitate.

Note : Slight precipitation in the medium may occur, which is inheritant property of the medium, and does not affect the performance of the medium.

### Principle And Interpretation

XLD Agar has been recommended for the identification of *Enterobacteriaceae* (7) and for the microbiological testing. XLD Agar was formulated by Taylor (1-5) for the isolation and differentiation of enteric pathogens including *Salmonella* Typhi from other *Salmonella* species of foods, water and dairy products (8-12). XLD Agar exhibits increased selectivity and sensitivity as compared to other plating media e.g. SS Agar (M108), EMB Agar (M022) and Bismuth Sulphite Agar (M027) (2, 4, 6, and 13-16). The media formulation does not allow the overgrowth of other organisms over *Salmonella* and *Shigella* (17). Samples suspected of containing enteric pathogens, along with other mixed flora, are initially enriched in Modified Semisolid RV Medium Base (M1482) (18). The medium contains yeast extract, which provides nitrogen and vitamins required for growth. Though the sugars xylose, lactose and sucrose provide sources of fermentable carbohydrates, xylose is mainly incorporated into the medium since it is not fermented by Shigellae but practically by all enterics. This helps in the differentiation of Shigella species. Sodium chloride maintains the osmotic balance of the medium. Lysine is included to differentiate the *Salmonella* group from the non-pathogens. Salmonellae rapidly ferment xylose and exhaust the supply. Subsequently lysine is decarboxylated by the enzyme lysine decarboxylase to form amines with reversion to an alkaline pH that mimics the *Shigella* reaction. However, to prevent this reaction by lysine-positive coliforms, lactose and sucrose are added to produce acid in excess. Degradation of xylose, lactose and sucrose to acid causes phenol red indicator to change its colour to yellow. Bacteria that decarboxylate lysine to cadaverine can be recognized by the appearance of a red colouration around the colonies due to an increase in pH. These reactions can proceed simultaneously or successively, and this may cause the pH indicator to exhibit various shades of colour or it may change its colour from yellow to red on prolonged incubation. To add to the differentiating ability of the formulation, an H<sub>2</sub>S indicator system, consisting of sodium thiosulphate and ferric ammonium citrate, is included for the visualization of hydrogen sulphide produced, resulting in the formation of colonies with black centers. The non-pathogenic H<sub>2</sub>S producers do not decarboxylase lysine; therefore, the acid reaction produced by them prevents the blackening of the colonies (1).

XLD Agar is both selective and differential medium. It utilizes sodium deoxycholate as the selective agent and therefore it is inhibitory to gram-positive microorganisms. Some *Proteus* strains may give red to yellow colouration with most colonies developing black centers, giving rise to false positive reactions. Non-enterics like *Pseudomonas* and *Providencia* may exhibit red colonies. *S.Paratyphi A*, *S.Choleraesuis*, *S.Pullorum* and *S.Gallinarum* may form red colonies without H<sub>2</sub>S, thus resembling *Shigella* species (19).

## Type of specimen

Clinical samples - Blood ; Food and dairy samples ; Water samples

## Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (20,21).

For food and dairy samples, follow appropriate techniques for sample collection and processing as per guidelines (8,10).

For water samples, follow appropriate techniques for sample collection, processing as per guidelines and local standards.(3)

After use, contaminated materials must be sterilized by autoclaving before discarding.

## Warning and Precautions :

In Vitro diagnostic Use only. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets

## Limitations :

Slight precipitation in the medium may occur, which is inheritant property of the medium, and does not affect the performance of the medium.

This medium is general purpose medium and may not support the growth of fastidious organisms.

Some *Proteus* strains may give red to yellow colouration with most colonies developing black centers, giving rise to false positive reactions.

Non-enterics like *Pseudomonas* and *Providencia* may exhibit red colonies.

*S.Paratyphi A*, *S.Choleraesuis*, *S.Pullorum* and *S.Gallinarum* may form red colonies without H<sub>2</sub>S, thus resembling *Shigella* species.

## Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

## Quality Control

### Appearance

Light yellow to pink homogeneous free flowing powder

### Gelling

Firm, comparable with 1.5% Agar gel

### Colour and Clarity of prepared medium

Red coloured clear to slightly opalescent gel forms in Petri plates

### Reaction

Reaction of 5.67% w/v aqueous solution at 25°C . pH : 7.4±0.2

### pH

7.20-7.60

### Cultural Response

Cultural response was observed after an incubation at 35-37°C for specified time. Recovery rate is considered as 100% for bacteria growth on Soyabean Casein Digest Agar.

### Cultural Response

Organism	Inoculum (CFU)	Growth	Observed Lot value (CFU)	Recovery	Colour of Colony	Incubation period
<i>Salmonella Typhimurium</i> ATCC 14028 (00031*)	50 -100	luxuriant	25 -100	>=50 %	red with black centres	18 -72 hrs

<i>Salmonella Abony</i> NCTC 6017 (00029*)	50 -100	good-luxuriant	25 -100	>=50 %	red with black centres	18 -72 hrs
<i>Escherichia coli</i> ATCC 8739 (00012*)	50 -100	fair	10 -30	20 -30 %	yellow	18 -72 hrs
<i>Escherichia coli</i> ATCC 25922 (00013*)	50 -100	fair	10 -30	20 -30 %	yellow	18 -72 hrs
<i>Escherichia coli</i> NCTC 9002	50 -100	fair	10 -30	20 -30 %	yellow	18 -72 hrs
<i>Proteus vulgaris</i> ATCC 13315	50 -100	good-luxuriant	25 -100	>=50 %	grey with black centres	18 -72 hrs
<i>Salmonella Paratyphi A</i> ATCC 9150	50 -100	good-luxuriant	25 -100	>=50 %	red	18 -72 hrs
<i>Salmonella Paratyphi B</i> ATCC 8759	50 -100	good-luxuriant	25 -100	>=50 %	red with black centres	18 -72 hrs
<i>Salmonella Enteritidis</i> ATCC 13076 (00030*)	50 -100	good-luxuriant	25 -100	>=50 %	red with black centres	18 -72 hrs
<i>Salmonella Typhi</i> ATCC 6539	50 -100	good-luxuriant	25 -100	>=50 %	red with black centres	18 -72 hrs
<i>Shigella dysenteriae</i> ATCC 13313	50 -100	good-luxuriant	25 -100	>=50 %	red	18 -72 hrs
<i>Shigella flexneri</i> ATCC 12022 (00126*)	50 -100	fair-good	15 -40	30 -40 %	red	18 -72 hrs
<i>Shigella sonnei</i> ATCC 25931	50 -100	fair-good	15 -40	30 -40 %	red	18 -72 hrs
# <i>Klebsiella aerogenes</i> ATCC 13048	50 -100	fair	10 -30	20 -30 %	yellow	18 -72 hrs
<i>Enterobacter cloacae</i> ATCC 13047 (00083*)	50 -100	fair	10 -30	20 -30 %	yellow	18 -72 hrs
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 25923 (00034*)	>=10 <sup>3</sup>	inhibited	0	0%		>=72 hrs
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 6538 (00032*)	>=10 <sup>3</sup>	inhibited	0	0%		>=72 hrs
<i>Enterococcus faecalis</i> ATCC 29212 (00087*)	>=10 <sup>3</sup>	inhibited	0	0%		>=72 hrs

Key : \*Corresponding WDCM numbers.

## Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 20-30°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Use before expiry date on the label.

Product performance is best if used within stated expiry period.

## Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (20,21).

**Please refer disclaimer Overleaf.**

## Reference

1. Taylor W. L., 1965, Am. J. Clin. Pathol., 44:471-475.
2. Taylor W. L. and Harris B., 1965, Am. J. Clin. Pathol., 44:476.
3. Taylor W. L. and Harris B., 1967, Am. J. Clin. Pathol., 48:350.
4. Taylor W. L. and Schelhart B., 1967, Am. J. Clin. Pathol., 48:356.
5. Taylor W. L. and Schelhart B., 1968, Am. J. Clin. Pathol., 16:1387.
6. Taylor W. L. and Schelhart B., 1969, Appl. Microbiol., 18:393-395.
7. Chadwick P., Delisle G. H and Byer M., 1974, Can. J. Microbiol., 20, 1653-1664.
8. Salfinger Y., and Tortorello M.L. Fifth (Ed.), 2015, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.
9. Baird R.B., Eaton A.D., and Rice E.W., (Eds.), 2015, Standard Methods for the Examination of Water and Wastewater, 23rd ed., APHA, Washington, D.C.
10. Wehr H. M. and Frank J. H., 2004, Standard Methods for the Microbiological Examination of Dairy Products, 17th Ed., APHA Inc., Washington, D.C.
11. Williams H., (Ed.), 2005, Official Methods of Analysis of the Association of Official Analytical Chemists, 19th Ed., AOAC, Washington, D.C.
12. FDA Bacteriological Analytical Manual, 2005, 18th Ed., AOAC, Washington, D.C.
13. Dunn C. and Martin W. J., 1971, Appl. Microbiol., 22, 17-22.
14. Rollender M. A., Beckford O., Belsky R. D and Kostroff B. 1969, Am. J. Clin. Pathol., 51, 284-286.
15. Taylor W. L. and Schelhart B., 1969, Appl. Micro. 18, 1387-1392.
16. MacCarthy M. D., 1966, N. Z. J. Med. Lab. Technol., 20, 127-131.
17. Isenberg H. D., Kominos S., and Sigeal M., 1969, Appl Microbiol., 18, 656-659.
18. Aspinall S. T., Hindle M. A. and Hutchinson D. N., 1992, Eur. J. Clin. Microbiol., Inf. Dis. 11, 936-939.
19. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. 1, Williams and Wilkins, Baltimore.
20. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2<sup>nd</sup> Edition.
21. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock, D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.


Revision : 02 / 2018



In vitro diagnostic medical device



CE Marking



Storage temperature



Do not use if package is damaged



HiMedia Laboratories Pvt. Limited,  
23 Vadhani Industrial Estate,  
LBS Marg, Mumbai-86, MS, India



CE Partner 4U, Esdoornlaan 13, 3951  
DB Maarn The Netherlands,  
[www.cepartner4u.eu](http://www.cepartner4u.eu)

**Disclaimer :**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.