



Technical Data

Luria Bertani Agar, Modified

M1151F

Luria Bertani Agar, Modified is used for the cultivation and maintenance of recombinant strains of *Escherichia coli* and may be used for routine cultivation of not particularly fastidious microorganisms.

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	10.000
Yeast extract	5.000
Sodium chloride	5.000
Agar	10.000
Final pH (at 25°C)	7.2±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 30 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Dispense as desired.

Principle And Interpretation

Luria Bertani Agar, Modified is prepared as described by Lennox (1) for cultivation and maintenance of recombinant strains of *Escherichia coli*. The medium is nutritionally rich for the growth of pure cultures of recombinant strains. Strains generally derived from *Escherichia coli K12* are deficient in Vitamin B synthesis which are further modified by specific mutation to create auxotrophic strains and are unable to grow on nutritionally deficient media.

Casein enzymic hydrolysate provides peptides and peptones while Vitamin B complex is provided by yeast extract. Sodium chloride provides sodium ions for the membrane transport and maintains osmotic equilibrium of the medium.

Quality Control

Appearance

Cream to yellow Homogeneous Free flowing powder

Gelling

Firm, comparable with 1.0% Agar gel.

Colour and Clarity of prepared medium

Yellow to amber Clear to slightly opalescent

Reaction

Reaction of 3.0% w/v aqueous solution at 25°C. pH : 7.2±0.2

pH

7.00-7.40

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Cultural Response

Organism	Inoculum (CFU)	Growth	Recovery
Cultural Response <i>Escherichia coli</i> ATCC 23724	50-100	luxuriant	≥70%
<i>Escherichia coli</i> ATCC 25922	50-100	luxuriant	≥70%

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

Reference

- 1.Lennox E.S., 1955, Transduction of Linked Genetic Characters of the host by bacteriophage P1., Virology, 1:190.
- 2.Atlas R.M., 1993, Handbook of Microbiological Media, Ed. by Parks L., CRC Press, Inc.

Revision : 2 / 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory,diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.