



HiCrome Candida Differential Agar Base

M1297AR

Intended use

HiCrome Candida Differential Agar Base is selective and differential medium for rapid isolation and identification of *Candida* species from mixed cultures from clinical and non-clinical samples

Composition**

Ingredients	Gms / Litre
Peptone	4.000
Chromogenic mixture	13.600
Agar	13.600
Final pH (at 25°C)	6.0±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 15.6 grams in 500 ml purified / distilled water. Add the rehydrated contents of one vial of HiCrome Candida Differential Selective Supplement (FD283R). Heat to boiling with frequent agitation to dissolve the medium completely. DO NOT AUTOCLAVE. Cool to 45-50°C. Mix well and pour into sterile Petri plates.

Principle And Interpretation

Perry and Miller (4) reported that *Candida albicans* produces an enzyme b -N-acetyl- galactosaminidase and according to Rousselle et al (5) incorporation of chromogenic or fluorogenic hexosaminidase substrates into the growth medium helps in identification of *C. albicans* isolates directly on primary isolation. HiCrome™ Candida Differential Agar Base incorporates two disphot X-NAG which detects the activity of hexosaminidase and BCIP which detects phosphatase activity.!

HiCrome™ Candida Differential Agar Base is a selective and differential medium, which facilitates rapid isolation of yeasts from mixed cultures and allows differentiation of *Candida* species namely *C. albicans*, *C. krusei*, *C. tropicalis* and *C. glabrata* on the basis of colouration and colony morphology. On this medium results are obtained within 48 hours and it is useful for the rapid and presumptive identification of common yeasts in Mycology and Clinical Microbiology Laboratory.

Peptone provides nitrogenous, carbonaceous compounds and other essential growth nutrients. Chloramphenicol from the supplement suppresses the accompanying bacterial flora. *C. albicans* appear as light green coloured smooth colonies, *C. tropicalis* appear as blue to metallic blue coloured raised colonies. *C. glabrata*, *C. kefir*, *C. parapsilosis* colonies appear as cream to white, beige/yellow due to natural pigmentation and some alkaline phosphatase activity, while *C. krusei* appear as pink-purple, fuzzy, dry colonies.

Type of specimen

Clinical samples - Blood ; Food and dairy samples

Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (2,3).

For food and dairy samples, follow appropriate techniques for sample collection and processing as per guidelines (1,6).

After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions :

In Vitro diagnostic Use only. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations :

1. Slight variation in colour for isolates may be observed as the reaction is based on the enzyme present in organism.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to beige homogeneous free flowing powder

Gelling

Firm, comparable with 1.36% Agar gel

Colour and Clarity of prepared medium

Light amber coloured, opaque gel forms in Petri plates

Reaction

Reaction of 3.12% w/v aqueous solution at 25°C. pH : 6.0±0.2

pH

5.80-6.20

Cultural Response

Cultural characteristics observed with added HiCrome Candida Differential Selective Supplement (FD283R) after an incubation at 20-25°C for 40-48 hours.

Cultural Response

Organism	Inoculum (CFU)	Growth	Recovery	Colour of Colony
Cultural Response				
<i>Candida albicans</i> ATCC 10231	50-100	good-luxuriant	≥50%	light green
<i>Candida krusei</i> ATCC 24408	50-100	good-luxuriant	≥50%	Purple, fuzzy
<i>Candida tropicalis</i> ATCC 750	50-100	good-luxuriant	≥50%	Blue to purple
<i>Candida kefyr</i> ATCC 66028	50-100	good-luxuriant	≥50%	Cream to white
<i>Candida parapsilosis</i> ATCC 22019	50-100	good-luxuriant	≥50%	Cream to white
<i>Candida glabrata</i> ATCC 15126	50-100	good-luxuriant	≥50%	Cream to white
<i>Escherichia coli</i> ATCC 25922 (00013*)	≥10 ³	inhibited	0%	
<i>Escherichia coli</i> ATCC 8739 (00012*)	≥10 ³	inhibited	0%	

Key : *Corresponding WDCM numbers.

Storage and Shelf Life

Store between 2-8°C in a tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Use before expiry date on the label.

Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (2,3).

Reference

1. American Public Health Association, Standard Methods for the Examination of Dairy Products, 1978, 14th Ed., Washington D.C.
2. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
3. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock, D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
4. Perry J. L. and Miller G. R., 1987, J. Clin. Microbiol., 25: 2424 -2425.
5. Rousselle P., Freydiere A., Couillerot P., de Montclos H. and Gille Y., 1994, J. Clin. Microbiol. 32:3034-3036.

6. Salfinger Y., and Tortorello M.L. Fifth (Ed.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.


Revision : 02/ 2018

 IVD

In vitro diagnostic medical device



CE Marking



Storage temperature



Do not use if package is damaged



HiMedia Laboratories Pvt. Limited,
23 Vadhani Industrial Estate,
LBS Marg, Mumbai-86, MS, India

 EC REP

CE Partner 4U ,Esdoornlaan 13, 3951
DB Maarn The Netherlands,
www.cepartner4u.eu

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.