



Brucella Agar Base w/ 1% Dextrose

M1638

Brucella Agar Base w/ 1% Dextrose is used for the cultivation and isolation of *Brucella* species.

Composition**

Ingredients	Gms / Litre
Peptic digest of animal tissue	10.000
Meat extract	5.000
Dextrose	10.000
Sodium chloride	5.000
Agar	15.000
Final pH (at 25°C)	7.5±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 22.5 grams in 500 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121oC) for 15 minutes. Cool to 45-50°C and aseptically add sterile 5% (v/v) inactivated horse serum RM1239 (Inactivate RM1239 by heating at 56°C for 30 minutes) and rehydrated contents of one vial Brucella Selective Supplement (FD005). Mix well before pouring into sterile Petri plates.

Principle And Interpretation

Brucellosis is a zoonotic disease with a domestic animal reservoir. It is an occupational disease of veterinarians, microbiologists, farmers etc. The route of infections is genital, nasopharyngeal, gastrointestinal, conjunctival, respiratory and through abraded skin (1, 2). Brucellosis in humans has a variable incubation period, an insidious or abrupt onset and no pathognomic symptoms or signs. Brucella Agar was designed for cultivating *Brucella* species from diagnostic specimens. With the incorporation of blood or other nutritious substances, it facilitates the cultivation of variety of fastidious anaerobic organisms (3). However, Brucella Medium is supplemented with antibiotics to prevent overgrowth of other accompanying organisms. Brucella Agar Base w/ 1.0% Dextrose was originally developed by Jones and Morgan (4) for preparations of serum-dextrose-antibiotic medium used for the isolation and cultivation of *Brucella* species.

The medium contains peptic digest of animal tissue and meat extract, which facilitates cultivation of variety of fastidious anaerobic organisms; by providing essential nutrients. Dextrose serves as source of energy. Addition of antibiotics (as FD) makes the medium highly selective for *Brucella* species. Ethyl violet and circulin, which were recommended initially, are no longer used (5).

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Light yellow coloured, clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 4.5% w/v aqueous solution at 25°C. pH : 7.5±0.2

pH

7.30-7.70

Cultural Response

M1638: Cultural characteristics observed in presence of 10% Carbon dioxide (CO₂) atmosphere with added 5% sterile inactivated horse serum and Brucella Selective Supplement (FD005), after an incubation at 35-37°C for 24-48 hours

Organism

Growth

Cultural Response

<i>Brucella melitensis</i> ATCC 4309	luxuriant
<i>Brucella suis</i> ATCC 4314	luxuriant
<i>Escherichia coli</i> ATCC 25922	inhibited
<i>Staphylococcus aureus</i> ATCC 25923	inhibited

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

1. Murray P. R., Baron E. J., Jorgensen J. H., Tenover F. C., Tenover F. C., (Eds.), 8th Ed., 2003, Manual of Clinical Microbiology, ASM, Washington, D.C.
2. Young E. J., 1983, Human Brucellosis, Rev. Infect. Dis., 5:821-842
3. Atlas R. M., 1997, Handbook of Microbiological Media, 2nd Ed., Parks L.C. (Ed.), CRC Press, New York.
4. Jones Lois M. and Brinley Morgan W. J., 1958, Bull. Wld. Hlth. Org., 19:200-203
5. Alton G. G. and Jones L. M., 1967, Lab Technique in Brucellosis, WHO, Geneva.

Revision : 1 / 2011

**Disclaimer :**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.