



HiCrome Malassezia Agar (Twin Pack)

M1985

Intended use

For isolation, cultivation and identification of *Malassezia furfur* from clinical samples.

Composition**

Ingredients	Gms / Litre
Part A	-
Peptone special	30.000
Chromogenic mixture	1.400
Agar	15.000
Part B	-
Tween 40	10.000
Glycerol	5.000
Final pH (at 25°C)	5.80±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 15ml of fluid Part B in 1000 ml distilled/purified water. Add 46.4 grams of Part A. Mix well and heat to boiling to dissolve the medium completely. DO NOT AUTOCLAVE. Cool to 45-50°C. Mix well and pour into sterile Petri plates .

Principle And Interpretation

Malassezia is a genus of fungi, naturally found on the skin surfaces of many animals, including humans. Media based on malt extract is appreciated by many microbiologists due to their richness and nutrient balance especially for the cultivation of fastidious microorganisms. With acidic pH, they are used for the isolation, cultivation and maintenance of yeast and moulds. *M. furfur* is a lipophilic yeast, therefore in vitro growth must be stimulated by natural oils or other fatty substances.

Peptone special provides nitrogenous and carbonaceous compounds, long chain amino acids, vitamins and other essential growth nutrients. Low pH favours fungal growth and inhibits contaminating bacteria from test samples (1). Tween 40, Glycerol enhances the growth of *Malessezia* species as it is a lipophilic yeast

Some pathogenic fungi may produce infective spores which are easily dispersed in air, so examination should be carried out in safety cabinet. For heavily contaminated samples, the plate must be supplemented with inhibitory agents for inhibiting bacterial growth with lower pH.

Type of specimen

Specimen Collection and Handling

2,3

Warning and Precautions :

For in vitro diagnostic use only. Read the label before opening the container. Wear protective gloves protective clothing eye protection face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets

Limitations :

Due to nutritional variation , some strains may show poor growth.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Part A : Cream to yellow homogeneous free flowing powder Part B: Colourless to pale yellow viscous solution

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Part A : Cream to yellow homogeneous free flowing powder Part B: Colourless to pale yellow viscous solution

Gelling

Firm, comparable with 1.5% Agar gel.

Colour and Clarity of prepared medium

Yellow coloured, opalescent gel with scum forms in Petri plates.

Reaction

Reaction of 4.64% w/v aqueous solution of Part A and 1.5% v/v of Part B at 25°C. pH : 5.80±0.2

pH

5.60-6.00

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 48-72 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony
<i>Malassezia furfur</i> ATCC 14521	50-100	good-luxuriant	≥50%	mauve, small
<i>Candida albicans</i> ATCC 10231	50-100	good-luxuriant	≥50%	pale green to green
<i>Candida glabrata</i> ATCC 15126	50-100	good-luxuriant	≥50%	colourless
<i>Candida krusei</i> ATCC 24408	50-100	good-luxuriant	≥50%	purple
<i>Candida tropicalis</i> ATCC 750	50-100	good-luxuriant	≥50%	metallic blue

Storage and Shelf Life

Store between 2-8°C in a tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Use before expiry date on the label.

Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (2,3).

Reference

- Murray PR, Baron EJ, Tenover JC, Tenover FC (editors) 2003, Manual of clinical Microbiology, 8th ed., ASM, Washington, D.C.
- Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
- Jorgensen, J.H., Tenover, J.C., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock, D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.


Revision : 01 / 2017



In vitro diagnostic medical device



CE Marking



Storage temperature



Do not use if package is damaged



HiMedia Laboratories Pvt. Limited,
B /4-6 , MIDC, Dindori, Nashik MH

www.himedialabs.com



CE Partner 4U ,Esdoornlaan 13, 3951
DB Maarn The Netherlands,

www.cepartner4u.eu

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.