



## Standard Nutrient Agar

M877

Standard Nutrient Agar is used as a general utility medium for cultivation and enumeration of not particularly fastidious microorganisms.

### Composition\*\*

Ingredients	Gms / Litre
Beef extract	10.000
Peptic digest of lean meat from	500.000
Sodium chloride	5.000
Agar	20.000
Final pH ( at 25°C)	7.6±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 45 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well before pouring in sterile Petri plates.

### Principle And Interpretation

Standard Nutrient Agar is formulated as per the recommendation of APHA as a general purpose medium for the cultivation of non-fastidious organisms from water and wastewater, dairy and food products (1, 2).

Peptic digest of lean meat provides the amino acids and large chain peptides. Beef extract (meat infusion) provides water soluble substances like carbohydrates, vitamins, organic nitrogen compounds and salts (3). Sodium chloride maintains osmotic equilibrium.

### Quality Control

#### Appearance

Yellowish brown coloured homogeneous free flowing powder

#### Gelling

Firm, comparable with 2.0% agar gel.

#### Colour and Clarity of prepared medium

Light amber coloured clear to slightly opalescent gel forms in petri plates.

#### Reaction

Reaction of 4.5% w/v aqueous solution at 25°C. pH : 7.6±0.2

#### pH

7.40-7.80

#### Cultural Response

M877: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours .

Organism	Inoculum (CFU)	Growth	Recovery
<i>Escherichia coli</i> ATCC 25922	50-100	Good - luxuriant	≥70%
<i>Staphylococcus aureus</i> ATCC 25923	50-100	Good - luxuriant	≥70%
<i>Pseudomonas aeruginosa</i> ATCC 27853	50-100	Good - luxuriant	≥70%
<i>Streptococcus pneumoniae</i> ATCC 6303	50-100	Good - luxuriant	≥70%
<i>Salmonella Typhi</i> ATCC 6539	50-100	Good - luxuriant	≥70%

## Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

## Reference

1. Greenberg A. E., Trussell R. R. and Clesceri L. S. (Eds.), 1985, Standard Methods for the Examination of Water and Wastewater, 16th ed., APHA, Washington, D.C.
2. Speck M. (Ed.), 1984, Compendium of Methods for the Microbiological Examination of Foods, 2nd ed., APHA, Washington, D.C.
3. Pelczar, Chan and Kreig, 1986, Microbiology, 5th ed., McGraw-Hill Book Company, New York.

Revision : 02 / 2015

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.