

MB602

HiPurA™ Total RNA Miniprep Purification Kit

Kit Contents

| Product Code | Reagents provided | MB602 | | |
|--------------|-----------------------------------------------------------------------|----------|----------|-------------|
| | | 20 Preps | 50 preps | 250 preps |
| DS0037 | RNA Lysis Solution (HRL) | 20 ml | 45 ml | 190 ml |
| DS0041 | Prewash Solution (RW1) | 20 ml | 45 ml | 190 ml |
| DS0012 | Wash Solution Concentrate (WS) | 7 ml | 15 ml | 65 ml |
| DS0042 | Elution Solution (RNase- Free Water) | 4 ml | 8 ml | 50 ml |
| DBCA03 | HiElute Miniprep Spin Column (Capped) [in DBCA016 Collection Tube] | 22 nos | 55 nos | 260 nos |
| DSCA02 | HiShredder [in DBCA016 Collection Tube] | 22 nos | 55 nos | 260 nos |
| DBCA016 | Collection Tube (Uncapped), Polypropylene (2.0 ml) | 20 nos | 50 nos | 250 nos |
| PW1139 | Collection Tube, Polypropylene (2.0 ml) | 40 nos | 100 nos | 2 X 250 nos |

Introduction

HiPurA Total RNA Miniprep Purification Kit provides a fast and easy method for purification of total RNA for Northern analysis, Poly A⁺ RNA selection, Primer extension, RNase and S1 nuclease protection assays, RT-PCR, Differential display, Expression-array and expression-chip analysis and cDNA library construction. The RNA purification procedure using the miniprep spin columns comprises of three steps viz, adsorption of RNA to the membrane, removal of residual contaminants and elution of pure RNA. HiMedia's HiElute Miniprep Spin column (Capped) format allows rapid processing of multiple samples. The columns have a high binding capacity and high quality RNA is obtained from various species. The RNA obtained is compatible with various downstream applications as mentioned above.

HiPurA™ Total RNA Miniprep Purification Kit

This kit simplifies isolation of RNA from animal cells, animal tissues and yeast, and for cleanup of RNA from crude enzymatic reactions (e.g. DNase digestion, Proteinase digestion, RNA ligation, and labeling reaction) with spin-column procedure. The lysis buffer provided in the kit helps in cell disruption and denaturation, samples are centrifuged through a HiShredder which removes insoluble material and reduces the viscosity of the lysate by disrupting viscous material. Ethanol is added to the cleared lysate, which promotes selective binding of RNA to the HiElute Miniprep Spin Column (Capped) membrane. After the initial binding of RNA, impurities like proteins, polysaccharides, low molecular weight metabolites and salts are removed by short washing steps. High quality RNA is finally eluted in the Elution Solution provided with the kit. A specialized high-salt buffer system allows upto 100 µg of total RNA longer than 200 bases to bind to the HiElute Miniprep Spin column (Capped) membrane.

HiElute Miniprep Spin Column (Capped) [DBCA03]

HiElute Miniprep Spin column (Capped) is based on the advanced silica binding principle presented in a microspin format. The system efficiently couples the reversible nucleic acid-binding properties of the advanced gel membrane and the speed plus versatility of spin

column technology to yield high quantity of RNA. The use of spin column facilitates the binding, washing, and elution steps thus enabling multiple samples to be processed simultaneously. This column eliminates the need for alcohol precipitation, expensive resins, and harmful organic compounds such as phenol and chloroform, otherwise employed in traditional RNA isolation techniques. RNA binds specifically to the advanced silica-gel membrane while contaminants pass through. PCR inhibitors such as divalent cations and proteins are completely removed in two efficient wash steps leaving pure nucleic acid to be eluted in the Elution Solution provided with the kit. The ratio of 28S rRNA to 18S rRNA should be 2:1. The ribosomal RNAs should appear as sharp bands or peaks.

Elution

The yield of RNA depends on the sample type and the number of cells in the sample. A single elution with 30-50 μ l of Elution Solution will provide sufficient RNA to carry out multiple amplification reaction.

Concentration, yield and purity of RNA

Spectrophotometric analysis and agarose gel electrophoresis will reveal the concentration and the purity of the RNA. Use Elution Solution to dilute samples and to calibrate the spectrophotometer, measure the absorbance at 260 nm, 280 nm and 320 nm using a quartz microcuvette. Absorbance readings at 260 nm should fall between 0.1 and 1.0. The 320 nm absorbance is used to correct for background absorbance. An absorbance of 1.0 at 260 nm corresponds to approximately 40 μ g/ml of RNA. The $A_{260}\text{-}A_{320}/A_{280}\text{-}A_{320}$ ratio should be 1.8 – 2.1. Purity is determined by calculating the ratio of absorbance at 260 nm to absorbance at 280 nm. RNA purified by HiPurA Total RNA Miniprep Purification Kit is free of protein and other contaminants that can inhibit PCR or other enzymatic reactions.

Concentration of RNA sample (μ g/ml) = 40 x A_{260} x dilution factor.

Materials needed but not provided

- 30°C water bath - For Isolation of total RNA from Yeast
- Tabletop Microcentrifuge (with rotor for 2.0 ml tubes)
- RNase – free pipette tips (aerosol barrier recommended)
- Ethanol (96 – 100%)
- Ethanol (70%)
- 2- mercaptoethanol (β -ME) (Product Code: MB041)
- Deoxyribonuclease I Solution (RNase-Free) and DNase Digest Buffer (procured from any standard company)
- Lyticase /zymolase (Product Code: MB099), 1 M Sorbitol and 0.1 M EDTA, pH 7.4 - For Isolation of total RNA from Yeast

Storage

HiPurA Total RNA Miniprep Purification Kit can be stored at room temperature (15-25°C) for up to 1 year without showing any reduction in performance.

Precautions to be taken while handling RNA

Ribonucleases (RNases) are very stable and active enzymes that generally do not require cofactors to function. Since RNases are difficult to inactivate and even minute amounts are sufficient to destroy RNA, do not use any plasticware or glassware without first eliminating possible RNase contamination. Great care should be taken to avoid inadvertently introducing RNases into the RNA sample during or after the isolation procedure. In order to create and maintain an RNase-free environment, the following precautions must be taken during pretreatment and use of disposable and non- disposable vessels and solutions while working with RNA.

1. Always wear latex or vinyl gloves while handling reagents and RNA samples to prevent RNase contamination from surface of the skin or from dusty laboratory equipment. Change gloves frequently and keep tubes closed whenever possible.
2. Use sterile, disposable plasticware and autoclavable pipettes reserved for RNA work to prevent cross-contamination with RNases from shared equipments.
3. Non-disposable plasticware should be treated before use to ensure that it is RNase-free. Plasticware should be thoroughly rinsed with 0.1M NaOH, 1mM EDTA followed by RNase-free water. Alternatively, chloroform-resistant plasticware can be rinsed with chloroform to inactivate RNases.
4. Glassware used for RNA work should be cleaned with a detergent, thoroughly rinsed, and oven baked at 240°C for four or more hours before use. Alternatively glassware can be treated with DEPC (Diethyl pyrocarbonate). Fill glassware with 0.1% DEPC (0.1% in water), allow to stand overnight at 37°C, and then autoclave or heat to 100°C for 15 min to eliminate residual DEPC.
5. Electrophoresis tanks should be cleaned with detergent solution (e.g., 0.5% SDS), thoroughly rinsed with RNase-free water, and then rinsed with ethanol and allowed to dry.
6. Solutions (water and other solutions) should be treated with 0.1% DEPC.

General Preparation Instructions

1. **β-mercaptoethanol (β-ME) must be added to RNA Lysis Solution (HRL) before use.**
β-ME is toxic; dispense in a fume hood and wear appropriate protective clothing. Add 10μl β-ME per 1ml Lysis Solution. Lysis Solution containing β-ME can be stored at room temperature for up to 1 month.
2. **Thoroughly mix reagents**
Examine the reagents for precipitation. If any kit reagent forms a precipitate (other than enzymes), warm at 55-65°C until the precipitate dissolves and allow cooling to room temperature (15-25°C) before use.
3. Ensure that clean & dry Nuclease-free tubes and tips are used for the procedure.
4. **Dilute Wash Solution Concentrate (WS) (DS0012) as follows:**

| Number of Preps | Wash Solution Concentrate (WS) | Ethanol (96-100 %) |
|-----------------|--------------------------------|--------------------|
| 20 | 7 ml | 21 ml |
| 50 | 15 ml | 45 ml |
| 250 | 65 ml | 195 ml |

5. **For Yeast protocol**

Buffer for Enzymatic lysis:

Depending on the yeast strain and enzyme used, the amount of enzyme and composition of this buffer may vary. Please adhere to guidelines of enzyme supplier. However, in most cases Buffer Y1 can be used.

Buffer Y1 1 M Sorbitol
 0.1 M EDTA, pH 7.4

Just before use, add:

10 μL of β-mercaptoethanol per 1 ml of Sorbitol buffer.

6. Preset the centrifuge at 4°C for Yeast RNA Isolation protocol.

Centrifugation

All centrifugation steps are carried out in conventional laboratory centrifuge e.g. Beckman CS-6KR, Heraeus Varifuge 3.0R, or Sigma 6k10 with fixed angle rotor. The tubes provided with the kit are compatible with almost all laboratory centrifuges and rotors. All centrifugation steps are performed at room temperature and are given in g; the correct rpm can be calculated using the formula:

$$RPM = \sqrt{RCF/1.118} \times 10^5 r$$

Where RCF = required gravitational acceleration (relative centrifugal force in units of g); r = radius of the rotor in cm; and RPM = the number of revolutions per minute required to achieve the necessary g-force.

Procedure

A. Protocol for Isolation of Total RNA from Animal Cells

1. Harvest the cells according to step 1a, 1b, and 1c

1a. Cells grown in suspension (do not use more than 1×10^7 cells):

Pellet up to 1×10^7 cells by centrifuging for 5 minutes at 300 x g (≈ 1500 rpm) in a Collection tube (not supplied). Discard the culture medium completely and continue with step 2.

NOTE: Incomplete removal of cell-culture medium will inhibit lysis and dilute the lysate, affecting the binding of RNA to the HiElute Miniprep Spin Column (Capped) membrane resulting in lower RNA yield.

1b. Cells grown in a monolayer (do not use more than 1×10^7 cells):

Direct lysis of the cells can be done in the cell-culture vessel (up to 10 cm diameter) or trypsinized and collected as a cell pellet prior to lysis. Cells grown in cell-culture flasks should always be trypsinized.

To lyse cells directly in culture dish:

Determine the number of cells. Completely aspirate the cell-culture medium, and proceed immediately to step 2.

NOTE: Incomplete removal of cell-culture medium will inhibit lysis and dilute the lysate, affecting the binding of RNA to the HiElute Miniprep Spin Column (Capped) membrane resulting in lower RNA yield.

To trypsinize and collect cells:

Determine the number of cells. Aspirate the medium, and wash the cells with PBS. Aspirate the PBS, and add 0.1–0.25% trypsin in PBS. After the cells detach from the dish or flask, add medium (containing serum to inactivate the trypsin), transfer the cells to an RNase-free glass or polypropylene centrifuge tube (not supplied), and centrifuge at 300 x g for 5 minutes. Completely aspirate the supernatant, and proceed to step 2.

NOTE: Incomplete removal of cell-culture medium will inhibit lysis and dilute the lysate, affecting the binding of RNA to the HiElute Miniprep Spin Column (Capped) membrane resulting in lower RNA yield.

2. Lysis reaction

- 2a. For pelleted cells, loosen the cell pellet thoroughly by flicking the tube. Add the appropriate volume of RNA Lysis Solution (HRL) (DS0037) (see the table below). Vortex or pipet to mix, and continue with step 3.

NOTE: Incomplete loosening of the cell pellet may lead to inefficient lysis and reduced RNA yields. Ensure that β -ME is added to RNA Lysis Solution (HRL) before use.

| Number of pelleted cells | RNA Lysis Solution (HRL) |
|---------------------------------|--------------------------|
| $<5 \times 10^6$ | 350 μ l |
| $5 \times 10^6 - 1 \times 10^7$ | 600 μ l |

- 2b. For Cells grown in a monolayer, add appropriate volume of RNA Lysis Solution (HRL) (see the table below) to the cell culture dish. Collect cell lysate into a collection tube (not supplied). Vortex or pipet to mix and ensure that no cell clumps are visible before proceeding to step 3.

NOTE: Regardless of the cell number, use the Lysis Solution volumes indicated to completely cover the surface of the dish. Ensure that β -ME is added to RNA Lysis Solution (HRL) before use.

| Dish diameter | RNA Lysis Solution (HRL) |
|---------------|--------------------------|
| < 6 cm | 350 μ l |
| 6-10 cm | 600 μ l |

3. Homogenize the lysate according to step 3a.

If processing $\leq 1 \times 10^5$ cells, homogenize by vortexing for 1 minute. After homogenization, proceed to step 4.

NOTE: Incomplete homogenization leads to significantly reduced RNA yields and can cause clogging of the HiElute Miniprep Spin Column(Capped). Homogenization with a HiShredder generally results in higher RNA yields.

- 3a. Pipet the lysate directly into a HiShredder (DSCA02) placed in a 2 ml uncapped collection tube, and centrifuge for 2 minutes at 13,000 X g (14,000 rpm). Prepare for binding step.

B. Protocol for Isolation of Total RNA from Animal Tissues

1. Fresh or frozen (up to 30 mg) tissue.

Immediately place the weighed (fresh or frozen) tissue in liquid nitrogen, and grind thoroughly with a mortar and pestle. Allow the liquid nitrogen to evaporate. Do not allow the samples to thaw. Proceed with the lysis.

2. Lysis reaction

Add appropriate volume of RNA Lysis Solution (HRL) (DS0037) (see the table below). Mix properly and transfer it to an RNase free 2.0 ml capped collection tube. Vortex or pipet to mix.

| Amount of starting material | RNA Lysis Solution (HRL) |
|-----------------------------|--------------------------|
| <20 mg | 350 μ l |
| 20 mg to 30 mg | 600 μ l |

NOTE: Incomplete loosening of the cell pellet may lead to inefficient lysis and reduced RNA yields. Ensure that β -ME is added to RNA Lysis Solution (HRL) before use.

3. Pipet the lysate directly into a HiShredder (DSCA02) placed in a 2 ml uncapped collection tube, and centrifuge for 2 min at full speed. Continue with binding.

C. Protocol for Isolation of Total RNA from Yeast

Important notes before starting

For RNA isolation from Yeast, cells should be harvested in log-phase growth. Use only freshly harvested cells for the protocol.

The Yeast culture should be harvested at 4°C. All subsequent steps of the protocol should be performed at room temperature (15-25°C).

Procedure

1. Harvest and Resuspend the Yeast cells

Harvest yeast cells by centrifugation at 1,000 X g for 5 minutes at 4°C. (Do not use more than 5×10^7 yeast cells). Remove the culture medium and discard.

2. Resuspend the cells in 600 μ l of freshly prepared Buffer Y1. Add 50U of zymolyase Or lyticase and incubate at 30°C for 30 minutes with gentle shaking to generate spheroplasts. **(Prepare Buffer Y1 as indicated in General Preparation Instructions)**

3. Centrifuge for 5 minutes at 6500 x g (10,000 rpm) to pellet spheroplasts. Carefully remove and discard the supernatant.

4. Lysis reaction

Add 350 μ l of RNA Lysis Solution (HRL) (DS0037) to the pellet. Vortex or pipet to mix.

NOTE: If insoluble material is visible, centrifuge for 2 minutes at maximum speed, and use only the supernatant in the subsequent steps. Ensure that β -ME is added to RNA Lysis Solution (HRL) before use.

5. Pipet the lysate directly into a HiShredder (DSCA02) placed in a 2 ml uncapped collection tube, and centrifuge for 2 minutes at 13,000 X g (14,000 rpm). Continue with binding.

D. Protocol for RNA Cleanup

Important notes before starting

A maximum of 100 μ g RNA can be used in the RNA cleanup protocol.

Procedure

1. Adjust sample to a volume of 100 μ l with RNase-free water. Add 350 μ l of RNA Lysis Solution (HRL) (DS0037) and mix thoroughly.

NOTE: Ensure that β -ME is added to RNA Lysis Solution (HRL) before use.

2. Prepare for binding:

Add 250 μ l of ethanol (96-100%) to the diluted RNA, and mix thoroughly by pipetting. Do not centrifuge. Continue without delay with column step (below) followed by washing with Wash Solution (WS) (DS0012) upto elution steps (below) [Skip the wash step with Prewash Solution (RW1) (DS0041) in RNA cleanup protocol].

RNA ISOLATION FROM ANIMAL CELLS, ANIMAL TISSUES, YEAST AND FOR RNA CLEANUP

• Prepare for binding:

Add 1 volume of 70% ethanol to the homogenized lysate, and mix well by pipetting. Do not centrifuge.

NOTE: A precipitate may be visible after addition of ethanol. This does not affect the procedure.

- **Load Lysate in HiElute Miniprep Spin Column (Capped) [DBCA03]**

Apply sample including any precipitate that may have formed, on the HiElute Miniprep Spin Column (Capped) . Close the tube gently, and centrifuge for 15 seconds at $\geq 8000 \times g$ ($\geq 10,000$ rpm). Discard the flow-through.

NOTE: If the volume exceeds 700 μl , load aliquots successively onto the HiElute Miniprep Spin column and centrifuge as above. Discard the flow-through after each centrifugation step.

Optional: On Column DNase digestion

Generally, DNase digestion is not required since the solutions of this kit efficiently remove most of the DNA without DNase treatment. However, further DNase treatment may be necessary for certain RNA applications that are sensitive to small amounts of DNA (e.g. TaqMan RT-PCR analysis with a low abundant target). DNA can also be removed by DNase digestion.

Carryout lysis, homogenization, and loading onto the HiElute Miniprep Spin Column (Capped) as indicated above. Instead of continuing with the Prewash Solution (RW1), follow steps 6a –6d below.

6a. Pipet 350 μl of Prewash Solution (RW1) (DS0041) into the HiElute Miniprep Spin column (Capped) , and centrifuge for 15 sec at $\geq 8000 \times g$ ($\geq 10,000$ rpm). Discard the flow through and reuse the collection tube in step 6c.

6b. Add 10 μl of DNase I Solution to 70 μl of DNase Digest Buffer. Mix by inversion. Do not vortex.

6c. Add 80 μl of DNase I/Digest Buffer mixture directly onto the HiElute Miniprep Spin Column (Capped). Incubate at room temperature for 15 minutes.

6d. Pipet 350 μl of Prewash Solution (RW1) into the HiElute Miniprep Spin column , and centrifuge for 15 seconds at centrifuge at $\geq 8000 \times g$ ($\geq 10,000$ rpm). Discard the flow-through and continue with the step 8.

Or

Alternatively, residual DNA can be removed by a DNase digestion after RNA isolation.

- **Prewash Solution (RW1)**

Add 700 μl of Prewash Solution (RW1) to the HiElute Miniprep Spin Column (Capped) centrifuge at $\geq 8000 \times g$ ($\geq 10,000$ rpm) for 1 minute. Discard the flow-through. Reuse the collection tube.

- Pipet 500 μl of Wash Solution (WS) (DS0012). Close the tube gently, and centrifuge for 1 minute at $\geq 8000 \times g$ ($\geq 10,000$ rpm) to wash the column. Discard the flow-through.

NOTE: Wash Solution (WS) is supplied as a concentrate. Ensure that ethanol is added to Wash Solution Concentrate (WS).

- Add another 500 μl of Wash Solution (WS) to the HiElute Miniprep Spin column (Capped). Close the tube gently, and centrifuge for 2 minutes at $\geq 8000 \times g$ ($\geq 10,000$ rpm) to dry the membrane. Continue directly with elution step or to eliminate any chance of possible Wash Solution carryover, perform the next step.

- **Optional:** Place the column in a new 2 ml collection tube (not supplied), and discard the old collection tube with the flow-through. Centrifuge at 13,000 X g (14,000 rpm) for 1 minute.
- **RNA Elution**
Transfer the HiElute Miniprep Spin column (Capped) to a new 2 ml capped collection tube. Pipet 30-50 µl Elution Solution (RNase-Free Water) directly onto the HiElute Miniprep Spin column (Capped). Close the tube gently, and centrifuge for 1 minute at ≥8000 x g (≥10,000 rpm) to elute.
- If the expected RNA yield is > 30 µg, repeat the elution step as described with a second volume of RNase-Free Water. Elute into the same collection tube.
- Transfer the eluate to a fresh capped 2ml collection tube for longer RNA storage.

NOTE: To obtain a higher total RNA concentration, this second elution step may be performed by using the first eluate. The yield will be 15-30% less than the yield obtained using a second volume of RNase-free water, but the final concentration will be higher.

Storage of the eluate with purified RNA: The eluate contains pure RNA, recommended to be stored at lower temperature (-80°C). Avoid repeated freezing and thawing of the sample which may cause denaturing of RNA.

References:

1. Sambrook, J., *et al.* Molecular Cloning: A laboratory Manual, 2nd ed. (Cold Spring Harbor Laboratory Press, Plainview, NY, 1989; pp. 7.3-7.5)
2. Farrell, Robert E., Jr.; RNA Methodologies; 2nd Edition; Academic Press: NY, 1998; pp. 37-53(Cat. No. Z350354)

Quality Control

Each lot of HiMedia’s HiPurA Total RNA Miniprep Purification Kit is tested against predetermined specifications to ensure consistent product quality.

Trouble shooting Guide:

| Sr. No. | Problem | Possible Cause | Solution |
|---------|-----------------------------------------------|-------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1. | Clogged HiElute Miniprep Spin Column (Capped) | Too much starting material | In subsequent preparations, reduce the amount of starting material. It is essential to use the correct amount of starting material (see protocols). |
| | | Tissues: Centrifugation before adding ethanol not performed | Centrifuge the lysate before adding ethanol and use only this supernatant in subsequent steps. Pellets contain cell debris can clog the HiElute Miniprep Spin Column (Capped). |
| | | Centrifugation temperature is too low | The centrifugation temperature should be 20 – 25°C. Some centrifuges may cool to below 20°C even when set at 20°C. This can cause formation of precipitates that can clog the column. If this happens, set the centrifugation temperature to 25°C. Warm the ethanol containing lysate to 37°C before transferring it to the column. |

| | | | |
|----|-----------------------------------------------------|------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 2. | Low RNA Yield | Too much of starting material | In subsequent preparations, reduce the amount of starting material. It is essential to use the correct amount of starting material (see protocols). |
| | | RNA still bound to HiElute Miniprep Spin Column | Repeat RNA elution, but incubate the column for 10 minutes at room temperature with Elution solution (RNase free water) before centrifuging. |
| | | Ethanol carryover | During the second wash with Wash Solution (WS) be sure to centrifuge at $\geq 8000 \times g$ ($\geq 10,000$ rpm) for 2 minutes to dry the column. After centrifugation, carefully remove the column from the collection tube so that the column does not contact the flow through otherwise carryover of ethanol will occur. To eliminate any chance of possible ethanol, centrifuge the column for another minute at full speed. |
| | | Incomplete removal of cell culture medium. | When processing cultured cells ensure complete removal of the cell culture medium after harvesting cells |
| 3. | Low A_{260} / A_{280} | Water has been used to dilute RNA for A_{260} / A_{280} measurement. | Use 10 mM Tris – Cl, pH 7.5, not RNase free water to dilute the sample before measuring purity. |
| 4. | DNA contamination in downstream experiments | No incubation with Prewash Solution (RW1) | In subsequent preparations, incubate the HiElute Miniprep Spin Column (Capped) for 5 minutes at room temperature after the addition of Prewash Solution (RW1) before centrifuging. |
| | | No DNase treatment | Follow the optional on-column DNase digestion step. |
| 5. | RNA does not perform well in downstream experiments | Ethanol carryover | During the second Wash using Wash Solution (WS), be sure to dry the HiElute Miniprep Spin Column membrane by centrifugation at $\geq 8000 \times g$ ($\geq 10,000$ rpm) for 2 minutes to dry the membrane. Following the centrifugation, remove the HiElute Miniprep Spin Column from the collection tube carefully so the column does not contact the flow-through as this will result in carryover of ethanol. |

Safety Information

The HiPurA Total RNA Miniprep Purification Kit is for laboratory use only, not for drug, household or other uses. Take appropriate laboratory safety measures and wear gloves when handling. Not compatible with disinfecting agents containing bleach. Please refer the Material Safety Data Sheet (MSDS) for information regarding hazards and safe handling practices.

Technical Assistance

At HiMedia we pride ourselves on the quality and availability of our technical support. For any kind of technical assistance mail to mb@himedialabs.com.

PIMB602_0/0514

MB602-07

Disclaimer :

User must ensure of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ Publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.