



L-Mono Differential HiCynth™ Agar Base

MCD1540

L. mono Differential HiCynth™ Agar Base has been recommended for the selective and differential isolation of *Listeria monocytogenes*.

Composition**

Ingredients	Gms / Litre
HiCynth™ Peptone No.1*	24.000
HiCynth™ Peptone No.5*	10.000
Sodium pyruvate	2.000
Glucose	2.000
Magnesium glycerophosphate	1.000
Magnesium sulphate	0.500
Sodium chloride	5.000
Lithium chloride	10.000
Disodium hydrogen phosphate anhydrous	2.500
Chromogenic substrate	0.050
Agar	15.000
Final pH (at 25°C)	7.2±0.2

**Formula adjusted, standardized to suit performance parameters

*Chemically defined peptones

Directions

Suspend 36.02 grams in 460 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Aseptically add sterile contents of 1 vial of L. mono Enrichment Supplement I (FD214) and sterile rehydrated contents of L. mono Selective Supplement I (FD212), L. mono Selective Supplement II (FD213). Mix well and pour into sterile Petri plates.

Warning : Lithium chloride is harmful. Avoid bodily contact and inhalation of vapours. On contact with skin wash with plenty of water immediately.

Principle And Interpretation

Listeria monocytogenes is a gram-positive foodborne human pathogen responsible for serious infections in pregnant women that may ultimately result in abortion, stillbirth, birth of a child with neonatal listeriosis and meningitis or primary bacteremia in adults and juveniles. The pathogenicity of *Listeria ivanovii* for humans is uncertain. Since *L. monocytogenes* and *L. innocua* have similar biochemical properties, they cannot be differentiated on traditional media (PALCAM, Oxford). L. mono Differential Agar Base is based on the formulation of Ottoviani and Agosti (1, 2) for the selective and differential isolation of *Listeria monocytogenes* from food and animal feeds which is adopted by ISO Committee (3). L. mono Differential HiCynth™ Agar Base is prepared by completely replacing animal or vegetable based peptones with chemically defined peptones to avoid BSE /TSE risks associated with animal peptones.

HiCynth™ peptone No.1 and HiCynth™ peptone No.5 provides nitrogenous and carbonaceous substances, long chain amino acids, vitamins and essential growth nutrients. Sodium pyruvate support good growth. Glucose is the fermentable carbohydrate. Sodium chloride maintains osmotic equilibrium. Phosphate buffers the medium. Lithium chloride and added selective supplements (FD212 and FD213) inhibit accompanying microflora and allow the growth of *Listeria* species. *Listeria* species hydrolyse the chromogenic substrate which produces green coloured colonies. Differentiation of *Listeria monocytogenes* from other *Listeria* species is based on phosphatidylinositol-specific phospholipase C (PIPLC) activity. Phospholipase C enzyme hydrolyses the purified substrate (FD214) added to the medium resulting in an opaque halo around *Listeria monocytogenes* colonies.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Please refer disclaimer Overleaf.

Colour and Clarity of prepared medium

Light amber coloured opalescent gel forms in Petri plates

Reaction

Reaction of 7.2% w/v aqueous solution at 25°C. pH : 7.2±0.2

pH

7.00-7.40

Cultural Response

Cultural characteristics observed with added sterile L. mono Selective supplement I (FD212), L. mono Selective Supplement II (FD213) and L.mono Enrichment supplement I (FD214) after an incubation at 35 - 37°C for 24 - 48 hours.

Cultural Response

Organism	Inoculum (CFU)	Growth	Recovery	Colour of Colony	PIPLC activity
Cultural Response					
<i>Candida albicans</i> ATCC 10231	≥10 ³	inhibited	0%		
<i>Enterococcus faecalis</i> ATCC 29212	≥10 ³	inhibited	0%		
<i>Escherichia coli</i> ATCC 25922	≥10 ³	inhibited	0%		
<i>Listeria innocua</i> ATCC 33090	50-100	luxuriant	≥50%	greenish-blue	negative
<i>Listeria grayi</i> ATCC 19120	50-100	luxuriant	≥50%	greenish-blue	negative
<i>Listeria ivanovii</i> ATCC 19119	50-100	luxuriant	≥50%	greenish-blue	positive, opaque halo around the colony exhibiting phosphatidylinositol specific phospholipase activity
<i>Listeria monocytogenes</i> ATCC 19112	50-100	luxuriant	≥50%	greenish-blue	positive, opaque halo around the colony exhibiting phosphatidylinositol specific phospholipase activity
<i>Listeria seeligeri</i> ATCC 35967	50-100	luxuriant	≥50%	greenish-blue	negative
<i>Listeria welshimeri</i> ATCC 43549	50-100	luxuriant	≥50%	greenish-blue	negative
<i>Pseudomonas aeruginosa</i> ATCC 27853	≥10 ³	inhibited	0%		

Storage and Shelf Life

Store dehydrated powder in tightly closed container and the prepared medium at 2-8° C. Use before expiry date on label.

Reference

- Ottaviani F., Ottaviani M., and Agosti M. (1997 a), Industrie Alimentari 36, 1-3.
- Ottaviani F., Ottaviani M., and Agosti M. (1997 b), Quimper Froid Symposium Proceedings p. 6, A.D.R.I.A. Quimper, France, 16-18 June 1997.
- Draft Amendment ISO 11290-2:1996/DAM

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.