



## Cetrimide Medium without Sterile Membrane Filter (Economy Pack)

MF007E

For detection and enumeration of *Pseudomonas* .

### Composition\*\*

Ingredients	Gms / Litre
Pancreatic digest of gelatin	20.000
Magnesium chloride	1.400
Potassium sulphate	10.000
Cetrimide	0.300

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

The test sample should be filtered through a sterile membrane filter having pore size of 0.22 $\mu$  / 0.45 $\mu$ . Rehydrate the nutrient pad with 2.0-2.5 ml sterile distilled / purified water. After filtration, remove the membrane filter aseptically using sterile forceps. Place the membrane filter on rehydrated nutrient pad. Incubate the inoculated nutrient. Interpret the results qualitatively by observing the presence or absence of growth and quantitatively by counting the number of colonies on the surface of the membrane filter and calculating CFU/ml.

### Principle And Interpretation

Field of Application: Water, pharmaceuticals, cosmetics and other samples. DriFilter Membrane Nutrient Pad Medium are ready to use sterile culture media in the form of a 50 mm biological inert absorbent pads impregnated with standard culture medium, then dried and sterilized in 55 mm petri plate. They eliminate the need of laborious media preparation and autoclaving procedures. The nutrient pads are to be just rewetted with sterile distilled water and are ready to use. Use of nutrient pads allows larger sample volumes to be tested at a time. Interpretation of results is directly by counting the CFUs and also quantifies the microbial load present in the sample. *Pseudomonas aeruginosa* grows well on all normal laboratory media but specific isolation of the organism, from environmental sites or from human, animal or plant sources, is best carried out on a medium, which contains a selective agent and also constituents to enhance pigment production. Most selective media depend upon the intrinsic resistance of the species to various antibacterial agents. Cetrimide inhibits the growth of many microorganisms whilst allowing *Pseudomonas aeruginosa* to develop typical colonies. Cetrimide is a quaternary ammonium salt, which acts as a cationic detergent that reduces surface tension in the point of contact and has precipitant, complexing and denaturing effects on bacterial membrane proteins. Cetrimide Agar developed by Lowburry (1) is a modification of Tech Agar (Medium A) with addition of 0.1% cetrimide for selective isolation of *P. aeruginosa* . Later, due to the availability of the highly purified cetrimide, its concentration in the medium was decreased (2). The incubation was carried out at 37°C for a period of 18-24 hours (3).

### Quality Control

#### Appearance

Dry filter membrane pad of 50mm diameter

#### Colour

Pale coloured nutrient pad

#### Sterility test

Passes release criteria

#### Cultural response

Cultural characteristics observed after incubation at 35-37°C for 18-24 hours

Organism	Growth	Colour of colony
<i>Ps.aeruginosa</i> ATCC 27853	Luxuriant	Colourless with blue green pigment

## Storage and Shelf Life

Store between 10-30°C. Use before expiry date on the label.

## Reference

- 1.King, Ward and Raney, 1954, J. Lab. Clin. Med., 44:301.
- 2.Lowbury, 1951, J. Clin. Pathol., 4:66.
- 3.Lowbury and Collins, 1955, J. Clin. Pathol., 8:47
- 4.Brown and Lowbury, 1965, J. Clin. Pathol., 18:752.
- 5.Murray P. R., Baron J. H., Pfaller M. A., Jorgensen J. H. and Tenover F. C., (Ed.), 2003, Manual of Clinical Microbiology, 8th Ed., American Society for Microbiology, Washington, D.C.
- 6.USFDA Bacteriological Analytical Manual, 2005, 18th Ed., AOAC, Washington, DC.
- 7.Forbes B. A., Sahm A. S. and Weissfeld D. F., Bailey & Scotts Diagnostic Microbiology, 10th Ed., 1998, Mosby, Inc., St. Louis, Mo.
- 8.Williams, (Ed.), 2005, Official Methods of Analysis of the Association of Official Analytical Chemists, 19th Ed., AOAC, Washington, D.C.
- 9.MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification -Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.
10. Goto and Enomoto, 1970, Jpn. J. Microbiol., 14:65.



### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.