



Chapman Stone Medium (without Membrane Filter) (Economy Pack)

MF008E

For detection and enumeration of *Staphylococci* .

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	10.000
Yeast extract	2.500
Gelatin	30.000
D-Mannitol	10.000
Sodium chloride	55.000
Ammonium sulphate	75.000
Dipotassium phosphate	5.000

**Formula adjusted, standardized to suit performance parameters

Directions

The test sample should be filtered through a sterile membrane filter having pore size of 0.22μ / 0.45μ . Rehydrate the nutrient pad with 2.0-2.5 ml sterile distilled / purified water. After filtration, remove the membrane filter aseptically using sterile forceps. Place the membrane filter on rehydrated nutrient pad. Incubate the inoculated nutrient. Interpret the results qualitatively by observing the presence or absence of growth and quantitatively by counting the number of colonies on the surface of the membrane filter and calculating CFU/ml.

Principle And Interpretation

Field of Application: Food, pharmaceuticals, cosmetics and other samples. DriFilter Membrane Nutrient Pad Medium are ready to use sterile culture media in the form of a 50 mm biological inert absorbent pads impregnated with standard culture medium, then dried and sterilized in 55 mm Petri plate. They eliminate the need of laborious media preparation and autoclaving procedures. The nutrient pads are to be just rewetted with sterile distilled water and are ready to use. Use of nutrient pads allows larger sample volumes to be tested at a time. Interpretation of results is directly by counting the CFUs and also quantifies the microbial load present in the sample. Chapman Stone medium is a selective media used for the isolation of food poisoning staphylococci. Foods commonly contaminated with *S. aureus* included synthetic creams, custards and high-salted food. Chapman Stone Agar is prepared according to the modification of Staphylococcus Medium 110 described by Chapman(1). It is similar to Staphylococcus Medium 110, previously described by Chapman (2), except that the sodium chloride concentration is reduced to 5.5% and additionally ammonium sulfate is included in the formulation. The main modification consist the inclusion of ammonium sulfate in the medium that allows the direct observation of gelatin hydrolysis, instead of adding reagents to the plate medium. Chapman Stone Medium is especially recommended for suspected food poisoning studies involving Staphylococcus (3). It is selective, due to the relatively high salt content, and is differential due to pigmentation, mannitol fermentation and the presence or absence of gelatin liquefaction. Material under test is inoculated on the surface and incubated at 30°C for 48 hours to produce separated colonies. After incubation, cream to golden yellow colonies are presumptively identified as *S. aureus* . While, white or non-pigmented colonies, are presumptively identified as *S. epidermidis* .

Quality Control

Appearance

Dry filter membrane pad of 50mm diameter

Colour

Pale coloured nutrient pad

Sterility test

Passes release criteria

Cultural response

Cultural characteristics observed after incubation at 35-37°C for 18-48 hours.

Organism	Growth	Colour of colony
<i>Staphylococcus epidermidis</i> ATCC 12228	Luxuriant	Colourless
<i>Staphylococcus aureus</i> ATCC 25923	Luxuriant	Colourless

Storage and Shelf Life

Store between 10-30°C. Use before expiry date on the label.

Reference

1. Chapman G. H., 1949, J. Bacteriol., 58:823 2. Chapman G. H., 1948, Food Res., 13:100. 3. Stone, 1935, Proc. Soc. Exp. Biol. N.Y., 33:18

**Disclaimer :**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.