



Antibiotic Assay Medium G

MM101

Antibiotic Assay Medium G is used for the microbiological assay of Amphotericin B and Nystatin using *Saccharomyces cerevisiae* as the test organisms in accordance with Indian Pharmacopoeia.

Composition**

Ingredients	Gms / Litre
Peptone	9.400
Yeast extract	4.700
Beef extract	2.400
Dextrose	10.000
Sodium chloride	10.000
Agar	23.500
pH after sterilization	6.1±0.1

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 60.0 grams in 1000 ml purified/distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Principle And Interpretation

The medium composition is in accordance to IP and CFR (1,2). This medium is used as seed agar for assay of antifungal agents like Amphotericin B and Nystatin. This medium is used for maintenance and inoculum development of *Saccharomyces cerevisiae*, indicator organism. This medium is also used for assaying mycostatic activity in pharmaceutical formulations. This medium is formulated as reported by Kirshbam and Arret (3)

Ingredients like peptone, yeast extract and beef extract supplements essential nutrients, minerals and growth factors for the growth of test organism. Dextrose in the medium provides enhanced source of carbon and energy. Osmotic equilibrium in the medium is maintained by sodium chloride which retains the cell integrity and viability.

Freshly prepared plates should be used for antibiotic assays. Test organisms are inoculated in sterile seed agar precooled to 40-45°C and spread evenly over the surface of solidified base agar. Prediffusion of antibiotics for 20 minutes in the agar by incubating at temperature below the optimal growth temperature for microorganism would facilitate better diffusion of antibiotic, followed by incubation of the plates for microbial growth.

Quality Control

Appearance

Cream to yellow coloured homogeneous free flowing powder

Gelling

Firm, comparable with 2.35% Agar gel.

Colour and Clarity of prepared medium

Yellow coloured clear to slightly opalescent gel forms in Petri plates.

pH

6.00-6.20

Cultural Response

MM101: Cultural characteristics observed after an incubation at 29-31°C for 48 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Antibiotics assayed
----------	----------------	--------	----------	---------------------

<i>Saccharomyces cerevisiae</i> ATCC 2601	50-100	luxuriant	$\geq 70\%$	Nystatin
<i>Saccharomyces cerevisiae</i> ATCC 9763	50-100	luxuriant	$\geq 70\%$	Amphotericin B

Storage and Shelf Life

Store below 30°C in tightly closed container and use freshly prepared medium . Use before expiry date on the label.

Reference

1. Indian Pharmacopoeia 2010, Ministry of Health and Family welfare, Government of India, New Delhi.
2. Tests and Methods of Assay of Antibiotics and Antibiotic containing Drugs, FDA, CFR, 1983 Title 21, Part 436, Subpart D, Washington, D.C : U.S. Government Printing Office, paragraphs 436, 100-436, 106, p. 242-259, (April).
3. Krishbam A and ArretB, 1967, J.Pharma. Sci. 56:512

Revision : 2 / 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.