

Dey-Engley Neutralizing Agar Plate (Gamma-irradiated)(Triple Pack)

Intended use

MP186GT

Dey-Engley Neutralizing Agar Plate is used in disinfectant testing, where neutralization of the chemical is important for determining its bactericidal activity.

Composition**

Ingredients	Gms / Litre
Tryptone	5.000
Yeast extract	2.500
Dextrose(Glucose)	10.000
Sodium thiosulphate	6.000
Sodium thioglycollate	1.000
Sodium bisulphite	2.500
Lecithin	7.000
Polysorbate 80	5.000
Bromocresol purple	0.020
Agar	15.000
Final pH (at 25°C)	7.6±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Either streak, inoculate or surface spread the test inoculum (50-100 CFU) aseptically on the plate.

Principle And Interpretation

Dey-Engley Neutralizing Agar Plate is formulated as per the procedure described by Engley and Dey (1). A strongly bacteriostatic substance inhibits the growth and reproduction of bacteria without killing them. These bacteria hold the ability to cause infection under favorable conditions. Dey-Engley Neutralizing Agar neutralizes a broad spectrum of antiseptics and disinfectants including quaternary ammonium compounds, phenolic, iodine and chlorine preparations, mercurial, formaldehyde and glutaraldehyde. (3).

Tryptone provide nitrogen and carbon source, long chain amino acids, vitamins and other essential nutrients. Dextrose is an energy source. Yeast extract is also a rich source of vitamin B-complex. The present formulation incorporate neutralizing substances for almost all the active products used as antiseptics and disinfectants. Sodium bisulfite neutralizes aldehydes; sodium thioglycollate neutralizes mercurial; sodium thiosulfate neutralizes iodine and chlorine (3); lecithin neutralizes quaternary ammonium compounds; and polysorbate 80, a non-ionic surface-active agent, neutralizes substituted phenolic (2,4,7,8). Bromocresol purple is an indicator for dextrose utilization. Due to the high concentration of lecithin in the broth medium, turbidity cannot be used to detect growth. Therefore, bromocresol purple and dextrose are added to the medium. Those organisms that ferment dextrose will turn the medium from purple to yellow. (3).

For Agar Medium: Dey -Engley Neutralizing Agar medium can be over-filled, producing a meniscus or dome-shaped surface that can be pressed onto a surface for sampling its microbial burden. Incubate the plates, by covering the lids, at an appropriate temperature. The presence of microorganism is determined by the appearance of colonies on the surface of agar medium. Neutralization Test: Growth in Neutralizing Broth and no growth in Neutralizing Broth Base indicate neutralization of disinfectant. To check bactericidal activity, both broth tubes are inoculated on D/E Neutralizing Agar. Positive growth from negative tubes of Neutralizing Broth Base indicates bacteriostatic substance while negative growth indicates a bactericidal disinfectant. All positive tubes should show growth on Dey-Engley Neutralizing Agar. The control disinfectants used in test procedure are 2% chlorine, 2% formaldehyde, 1% glutaraldehyde, 2% iodine, 2% phenol, 1/750 quaternary ammonium compounds, 1/1000 mercurial etc.

Type of specimen

Food and dairy samples; Environmental samples.

Specimen Collection and Handling

For food and dairy samples, follow appropriate techniques for sample collection and processing as per guidelines (1,3,8).

For water samples, follow appropriate techniques for sample collection, processing as per guidelines and local standards.(7)

After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions :

Read the label before opening the pack. Wear protective gloves/protective clothing/eye protection/ face protection.

Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations :

1. Due to nutritional variations, some strains may show poor growth

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Sterile Dey-Engley Neutralizing Agar in 90 mm disposable plates.

Colour of medium

Purple to reddish purple coloured medium

Quantity of medium

30 ml of medium in 90 mm disposable plates.

pH

7.40-7.80

Dose of irradiation (Kgy)

13.00- 20.00

Sterility Test

Passes release criteria

Cultural Response

Cultural characteristics observed after an incubation at 35 - 37°C for 40 - 48 hours.

Organism	Inoculum (CFU)	Growth	Recovery
<i>Escherichia coli</i> ATCC 25922 (00013*)	50-100	luxuriant	≥70%
<i>Pseudomonas aeruginosa</i> ATCC 27853 (00025*)	50-100	luxuriant	≥70%
<i>Salmonella</i> Typhimurium ATCC 14028 (00031*)	50-100	luxuriant	≥70%
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 25923 (00034*)	50-100	luxuriant	≥70%
<i>Bacillus subtilis</i> subsp. <i>spizizenii</i> ATCC 6633 (00003*)	50-100	luxuriant	≥70%
<i>Candida albicans</i> ATCC 10231 (00054*)	50-100	luxuriant	≥70%
<i>Aspergillus brasiliensis</i> ATCC 16404 (00053*)	50-100	luxuriant	≥50%

* - Corresponding WDCM numbers

Storage and Shelf Life

On receipt store between 20-30°C Use before expiry date on the label.
Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (5,6).

Reference

1. American Public Health Association, Standard Methods for the Examination of Dairy Products, 1978, 14th Ed., Washington D.C.
2. Brummer B., 1976, Appl. Environ. Microbiol., 32:80.
3. Engley and Dey, 1970. Chem. Spec. Manuf. Assoc. Proc., Mid-Year Meet., p. 100.
4. Erlandson A. L., and Lawrence C. A., 1953, Science 118:274.
5. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
6. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock, D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
7. Quisno R.A., Gibby I.W., and Foter M.J., 1946, Am. J. Phar., 118:320.
8. Salfinger Y., and Tortorello M.L. Fifth (Ed.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.
9. Wehr H. M. and Frank J. H., 2004, Standard Methods for the Microbiological Examination of Dairy Products, 17th Ed., APHA Inc., Washington, D.C.

Revision : 01 / 2019

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.