



## Tryptone T, Type-III

RM029

### Principle And Interpretation

Tryptone T Type III is manufactured under controlled conditions by enzymic hydrolysis. It is used for production of Tetanus toxin. It is also used as media ingredient for the cultivation of bacteria, fungi, moulds and yeasts.

### Quality Control

#### Appearance

Off white to light yellow homogenous free flowing powder, having characteristic but not putrescent odour.

#### Solubility

Freely soluble in distilled/ purified water, insoluble in alcohol.

#### Clarity

1% w/v aqueous solution is clear to slight opalescent after autoclaving at 15 lbs pressure (121°C) for 15 minutes.

#### Reaction

Reaction of 2% w/v aqueous solution at 25°C.

#### pH

5.50 - 7.50

#### Microbial Load:

##### Total aerobic microbial count (cfu/gm)

By plate method when incubated at 30-35°C for not less than 3 days.

Bacterial Count :  $\leq$  2000 CFU/gram

##### Total Yeast and mould count (cfu/gm)

By plate method when incubated at 20-25°C for not less than 5 days.

Yeast & mould Count :  $\leq$  100 CFU/gram

#### Test for Pathogens

1. *Escherichia coli*- Negative in 10 gms of sample 2. *Salmonella* species- Negative in 10 gms of sample 3. *Pseudomonas aeruginosa*- Negative in 10 gms of sample 4. *Staphylococcus aureus*- Negative in 10 gms of sample 5. *Candida albicans*- Negative in 10 gms of sample 6. *Clostridia*- Negative in 10 gms of sample

#### Indole test

Tryptophan content: Passes

#### Cultural response

Cultural response observed after an incubation at 35-37°C for 18-24 hours by preparing Tryptone Broth (M463) using Tryptone T Type III as an ingredient.

#### Cultural response

Organism	Growth	Indole reaction
<i>Escherichia coli</i> ATCC 25922 (WDCM 00013)	Luxuriant	Positive reaction, red ring at the interface of the medium
* <i>Klebsiella aerogenes</i> ATCC 13048 (WDCM 00175)	Luxuriant	Negative reaction, no colour development /cloudy ring
<i>Klebsiella pneumoniae</i> ATCC 13883(WDCM00097)	Luxuriant	Negative reaction, no colour development /cloudy ring

\*Formerly known as *Enterobacter aerogenes*

**Chemical Analysis**

Total Nitrogen	$\geq 12.0\%$
AminoNitrogen	$\geq 3.50\%$
Sodium chloride	$\leq 4.0\%$
Loss on drying	$\leq 5.0\%$
Residue on ignition	$\leq 15\%$
Iron (Fe)	$\leq 0.003\%$

**Storage and Shelf Life**

Store between 10- 30°C in tightly closed container and away from bright light. Use before expiry date on label. On opening, product should be properly stored in dry ventilated area protected from extremes of temperature and sources. Seal the container tightly after use.

Revision : 08 / 2021

**Disclaimer :**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.